

# Scoped Environmental Impact Study 10249 Hunsden Sideroad

Caledon, Ontario

#### Submitted to:

Carringwood Homes 101 Regent St Richmond Hill, ON L4C 9P4

#### Submitted by:

GEI Consultants Ltd. 1266 South Service Road, Unit C3-1 Stoney Creek, Ontario L8E 5R9 1-800-810-3281

April 2024 Project 2101948

#### **Statement of Conditions**

This Report / Study (the "Work") has been prepared at the request of, and for the exclusive use of, the Owner / Client, the Town of Caledon, Regional Municipality of Peel, Nottawasaga Valley Conservation Authority, and its affiliates (the "Intended User"). No one other than the Intended User has the right to use and rely on the Work without first obtaining the written authorization of GEI Consultants Ltd and the Owner. GEI Consultants Ltd. expressly excludes liability to any party except the Intended User for any use of, and/or reliance upon, the work.

Neither possession of the Work, nor a copy of it, carries the right of publication. All copyright in the Work is reserved to GEI Consultants Ltd. The Work shall not be disclosed, produced or reproduced, quoted from, or referred to, in whole or in part, or published in any manner, without the express written consent of GEI Consultants Ltd., the Town of Caledon, Regional Municipality of Peel, Nottawasaga Valley Conservation Authority, or the Owner.

# **Table of Contents**

1.	intro	auction	1				
	NI. 4	and the state of t					
2.		ral Heritage Planning Considerations	2				
	2.1	Town of Caledon Official Plan (TCOP)	2				
	2.2	Region of Peel Official Plan	4				
		2.3 Oak Ridges Moraine Conservation Plan					
	<ul><li>2.4 Provincial Policy Statement</li><li>2.5 Nottawasaga Valley Conservation Authority</li></ul>						
	2.5 2.6	6 6					
3.	Data	Collection and Analyses	7				
	3.1 Background References						
		3.1.1 Land Information Ontario	7 7				
		3.1.2 Natural Heritage Information Centre	7 7 8				
		3.1.3 Ontario Breeding Bird Atlas	8				
		3.1.4 Ontario Reptile and Amphibian Atlas	8				
		3.1.5 Ontario Butterfly and Moth Atlas	9				
		3.1.6 Citizen Science Database (iNaturalist)	9				
		3.1.7 Citizen Science Database (eBird)	9				
		3.1.8 Aquatic Species at Risk Distribution Mapping	9				
		3.1.9 Innisfil Creek Subwatershed Health Check 2023	10				
	3.2	Technical Methods and Field Studies	10				
		3.2.1 Vegetation and ELC Methods	10				
		3.2.2 Headwater Drainage Feature Assessment	11				
		3.2.3 Breeding Bird Surveys	11				
4.	Bio-Physical Characterization						
	4.1	Physiography	12				
	4.2	Landscape Ecology	12				
	4.3	Vegetation	12				
		4.3.1 Ecological Land Classification	12				
	4.4	4.3.2 Vascular Plants	13				
	4.4	Wildlife	14				
		4.4.1 Breeding Bird Surveys	14				
	4.5	4.4.2 Incidental Wildlife Headwater Drainage Features	15 15				
	4.5	neadwater Drainage Features	13				
5.	Analysis Of Ecological and Natural Heritage Significance						
	5.1	Wetlands/Significant Wetlands	17				
	5.2	Significant Coastal Wetlands	17				
	5.3	Habitat of Endangered and Threatened Species	18				
		5.3.1 Butternut	18				
	ΕΛ	5.3.2 Endangered Species of Bats Fish Habitat	18 18				
	5.4 5.5		19				
	5.6	Significant Woodlands/Region of Peel Woodlands Significant Valleylands	20				
	5.0	Olyminicant valicylatius	20				



9.	Conci	usions and Recommendations	36		
٥	Const	usions and Basemmandations	26		
8.	Enviro	onmental Management/Reforestation Plan	33		
	7.6	Air Pollution	32		
	7.5	Migratory Birds/Endangered Species of Bats	32		
	7.4	PNAC Woodland	32		
	7.3	Significant Woodlands/Habitat of Endangered and Threatened Species/Confirmed and Candidate Significant Wildlife Habitat	31		
		7.2.2.1 Stormwater Management and Changes in Infiltration 7.2.2.2 Impacts on Water Quality	30 31		
		7.2.2 Potential Post-Construction Impacts	30		
		7.2.1 Potential Impacts During Construction	28		
	7.2	Indirect Fish Habitat	28		
		7.1.3 Linkages	28		
		7.1.2 PNAC Woodland	27		
		7.1.1 Significant Woodlands	26		
<u>' ·                                     </u>	7.1	Vegetation Protection Zone	26		
7.	Imnac	t Assessment, Mitigation and Enhancement Opportunities	26		
6.	Description Of Proposed Development				
	5.15	Summary of Ecological Components Subject to Impact Assessment	24		
	5.14	EZ Designations	23		
	5.13	1 0 1 0			
	5.12		22		
	5.11	Kettle Lakes	22		
	5.10	Permanent and Intermittent Streams	22		
	5.6 5.9	Sand Barrens, Savannahs and Tallgrass Prairies	20 22		
	5.7 Significant Areas of Natural and Scientific Interest (ANSIS) 5.8 Significant Wildlife Habitat				
	5.7	Significant Areas of Natural and Scientific Interest (ANSIs)	20		

# Appendices A. Figures B. Tables

- C. Draft Plan

## 1. Introduction

GEI Consultants (GEI) was retained by Carringwood Homes to prepare an Environmental Impact Study (EIS) in support of a proposed residential development on lands known as 10249 Hunsden Sideroad (herein referred to as the Subject Lands), within the Town of Caledon, Ontario (**Figure 1, Appendix A**). The Subject Lands are legally known as Part Lots 25 and 26, Concession 9 Albion.

The Subject Lands presently consist predominantly of row-crop agricultural lands, currently planted in soy, with surrounding mature woodland communities within the southern and northeastern extents of the Subject Lands. The Subject Lands are situated within the Palgrave Estates Residential Community Secondary Plan area within the Town of Caledon, ON. The surrounding lands consist of a mixture of agricultural lands to the north, continuations of the natural areas present on the Subject Lands, a recently developed residential subdivision to the south, and rural residential lots.

A scoped EIS is required to assess the potential impacts of the proposed development on the natural heritage features and associated ecological functions on the Subject Lands. This report provides a characterization of the existing natural heritage conditions of the Subject Lands based on ecological investigations completed in 2021 and 2022.

This work considers applicable provincial and municipal requirements and policies including reference to the natural heritage policies of the Province of Ontario's Provincial Policy Statement (PPS; MMAH 2020) and associated provincial implementation guidance contained in the Natural Heritage Reference Manual (NHRM; MNR 2010).

This scoped EIS is a requirement of the municipal planning process and is intended to address the policies of the Town of Caledon, Region of Peel, and the Nottawasaga Valley Conservation Authority (NVCA).

The study components to date, have included:

- A review of existing background information, policies and legislation applicable to the Subject Lands in its regional context;
- A field review and description of the natural environmental features and functions on, and immediately adjacent to, the Subject Lands through the completion of various ecological surveys and inventories;
- Identification and delineation of key natural heritage features (KNHF) and key hydrologic features (KHF) in accordance with the ORMCP;
- An evaluation of the sensitivity of the natural heritage features and their functions on the Subject Lands;
- An assessment of linkage functions between the identified natural heritage features;
- A description of the proposed development based on the Draft Plan;
- Identification and discussion of the impacts that could affect the natural heritage features as a result of the proposed development; and
- Recommendations for mitigation to avoid or minimize impacts.



# 2. Natural Heritage Planning Considerations

The Subject Lands are subject to federal, provincial, and municipal legislation as well as land use policies established by the Town of Caledon, and the NVCA.

An assessment of the quality and extent of natural heritage features found on and adjacent to the Subject Lands was completed. Ecological opportunities and constraints to development were evaluated in the context of the requirements of the following regulatory agencies, local and regional municipalities, and/or legislation:

- Town of Caledon Official Plan (TCOP; 2018 Office Consolidation);
- Region of Peel Official Plan (ROP; 2022);
- Oak Ridges Moraine Conservation Plan (ORMCP; 2017)
- Provincial Policy Statement (PPS; 2020);
- Nottawasaga Valley Conservation Authority (NVCA); and
- Provincial Endangered Species Act, 2007 (ESA; 2007).

The relevant portions of each of these, as they apply to the Subject Lands and the development potential, are discussed in the following sections.

## 2.1 Town of Caledon Official Plan (TCOP)

The Subject Lands are situated within the limits of the Palgrave Estate Residential Community Secondary Plan Area. The secondary plan for this area provides protection from development to woodlots, wetlands and other ecologically significant areas, including valley and stream corridors (Policy 7.1.2.3), ORMCP Key Natural Heritage Features (KNHF) and Hydrologically Sensitive Features (also referred to as key hydrologic features, KHF) (Policy 7.1.2.12), and ORMCP Natural Core and Natural Linkage Areas (Policy 7.1.2.13).

Schedule G of the TCOP shows the Subject Lands situated within a mixture of Policy Areas 2, 3 and 4. Per TCOP Policies within section 7.1.5, Policy Areas 2 and 3 correspond with ORMCP Countryside Areas and are suitable for estate residential development, while Policy Area 4 corresponds with ORMCP Natural Core and Natural Linkage Areas and is unsuitable for estate residential development.

Schedule I of the TCOP shows the Subject Lands contain settlement area with Environmental Zone 1 (EZ1) and Environmental Zone 2 (EZ2) Designations. Per TCOP Policy 7.1.9, EZ1 designations includes all ORMCP KNHF and KHF, and their related Minimum Vegetation Protection Zones (MVPZ), along with more sensitive biological communities, valley and stream corridors and their associated floodplains, native upland and lowland woodlands, natural waterbodies, Provincially and locally significant wetlands, and Environmentally Significant/ Sensitive Areas along with other features of local or regional importance. EZ2 Designations are locations with high groundwater table, seasonal flooding, dry swale lowlands and natural depressions performing natural run-off, detention and groundwater recharge functions, and smaller hedgerows and strips of native vegetation. Though general mapping is provided within Schedule I, the actual limits of the features are to be determined through detailed studies, including Natural Heritage Evaluations (NHE) and/or Hydrological Evaluations.

The TCOP notes that development is not permitted within areas designated as EZ1, while limited development may be permitted within areas designated as EZ2, such as crossing of a narrow point of EZ2 with a driveway to permit reasonable access to a development lot.

## 2.2 Region of Peel Official Plan

The ROP implements the PPS natural features policies through the Greenlands System's Core Areas, Natural Areas and Corridors (NAC) and Potential Natural Areas and Corridors (PNAC) policy framework. The ROP natural heritage policies and identifies the following components as Core Areas (Schedule A) of the Peel Greenlands system (section 2.14.12):

- Significant wetlands;
- Significant coastal wetlands;
- Core woodlands meeting one or more criteria in Table 1 (of the ROP);
- Environmentally Sensitive or Significant Areas;
- Provincial Life Science Areas of Natural and Scientific Interest (ANSIs);
- Significant habitats of threatened and endangered species;
- Escarpment Naturals Areas of the Niagara Escarpment Plan; and
- Core Valley and Stream corridors meeting one or more criteria in Table 2 (of the ROP).

The ROP further identifies the following components as NAC (2.14.18):

- Evaluated non-provincially significant wetlands and coastal wetlands;
- Woodlands meeting one or more of the criteria for NAC woodland in Table 1 (of the ROP);
- Significant wildlife habitat;
- fish habitat;
- habitat of aquatic species at risk;
- habitat of endangered and threatened species defined in accordance with the Endangered Species Act;
- regionally significant life science ANSIs;
- provincially significant earth science ANSIs;
- Escarpment Protection Areas of the Niagara Escarpment Plan;
- the Lake Ontario shoreline and littoral zone and other natural lakes and their shorelines;
- any other valley and stream corridors that have not been defined as part of the Core Areas;
- sensitive headwater areas and sensitive ground water discharge areas; and
- any other natural features and functional areas interpreted as part of the Greenlands System NAC by the local municipalities, in consultation with the conservation authorities and the Ministry of Northern Development, Mines, Natural Resources and Forestry, including, as appropriate, elements of the PNAC.

PNAC are defined in the ROP as the following components (2.14.19):

- unevaluated wetlands and coastal wetlands;
- cultural woodlands and cultural savannahs within the Urban System meeting one or more of the criteria for PNAC woodland in Table 1 (of the ROP). The evaluation of cultural woodlands and cultural savannahs is also subject to Policy 2.14.29 of the ROP;

- any other woodlands greater than 0.5 hectares;
- regionally significant earth science ANSIs;
- sensitive ground water recharge areas;
- portions of Historic shorelines;
- open space portions of the Parkway Belt West Plan Area;
- · enhancement areas, buffers and linkages; and
- any other natural features and functional areas interpreted as part of the Greenlands System PNAC, by the individual local municipalities in consultation with the conservation authorities.

Schedule C2 of the ROP identifies the woodland situated at the southern extent of the Subject Lands as Core Area of the Regional Greenlands System, while Schedule C1 shows the limits of the Regional Greenlands System extending north to an east-west corridor along Hunsden Sideroad. Figure 7 of the ROP shows NAC associated with the woodlands in the northeast corner of the Subject Lands, and PNAC associated with the woodland west of the Subject Lands. The remaining areas of the Subject Lands are identified as within the limits of the Palgrave Estate Residential Community.

## 2.3 Oak Ridges Moraine Conservation Plan

The Subject Lands occur within the Oak Ridges Moraine physiographic region. The Subject Lands contain a mixture of Natural Linkage Area as well as Countryside Areas within the Palgrave Estate Residential Community.

The purpose of Natural Linkage Areas is to protect critical natural and open space linkages between the Natural Core Areas and along rivers and streams and maintain and where possible improve or restore the ecological integrity of the Plan Area. Applications must ensure connectivity between KNHF and KHF is maintained within and adjacent to the Natural Linkage Areas.

Residential development is expressly permitted within Palgrave Estates Residential Community of the Countryside Area of the ORMCP subject to the requirements of the Town of Caledon Official Plan, and various sections within the ORMCP.

A NHE and a hydrological evaluation are required with respect to the development of land within the minimum area of influence, but outside the related MVPZ of a KNHF or KHF, respectively.

The ORMCP defines KNHF and KHF and stipulates where development is or is not permitted.

KNHF are defined in Section 22(1) as one or more of the following:

- Wetlands;
- Habitat of endangered and threatened species;
- Fish habitat;
- ANSI (life science);



- Significant valleylands;
- Significant woodlands;
- Significant wildlife habitat (including habitat of special concern species); and/or
- Sand barrens, savannahs and tallgrass prairies.

KHF are defined in Section 26(1) as:

- Permanent and intermittent streams;
- Wetlands:
- Kettle Lakes; and
- Seepage areas and springs.

## 2.4 Provincial Policy Statement

The PPS (2020) provides direction on matters of provincial interest related to land use planning and development. It "...supports a comprehensive, integrated and long-term approach to planning..."

The PPS is to be read in its entirety and land use planners and decision-makers need to consider all relevant policies and how they work together.

Policies in section 2.1 (Natural Heritage) of the PPS identify eight types of significant natural heritage features, as follows:

- Significant wetlands;
- Significant coastal wetlands;
- Significant woodlands;
- Significant valleylands;
- Significant wildlife habitat (SWH);
- Fish habitat:
- Habitat of endangered and threatened species; and
- ANSIs.

Development and site alteration shall not be permitted in significant wetlands or significant coastal wetlands. Development and site alteration shall not be permitted in significant woodlands, significant valleylands, SWH or significant ANSIs, unless it is demonstrated that there will be no negative impacts on the natural features or their ecological functions.

Development and site alteration shall not be permitted in the habitat of endangered and threatened species or in fish habitat, except in accordance with provincial and federal requirements. Development and site alteration may be permitted on lands adjacent to fish habitat provided it has been demonstrated that there will be no negative impacts on the natural feature or their ecological functions.

## 2.5 Nottawasaga Valley Conservation Authority

The NVCA conducts reviews of planning processes associated with future development of properties within its jurisdictional boundaries. In addition, the NVCA provides planning and technical advice to planning authorities to assist them in fulfilling their responsibilities regarding natural hazards and wetlands and other relevant policy areas pursuant to the Planning Act as a watershed-based resource management agency, in addition to their Regulatory responsibilities. The Regulation Limit delineates hazardous lands, wetlands, shorelines and areas susceptible to flooding and associated allowances.

Pursuant to the Development, Interference with Wetland and Alterations to Shorelines and Watercourse Regulation (NVCA; Ontario Regulation 172/06), any development in or on areas defined in the Regulation (e.g., river or stream valleys, hazardous land, wetlands) requires permission from the Conservation Authority. The Conservation Authority may grant permission for development in or on these areas if, in its opinion, the control of flooding, erosion, dynamic beaches, pollution or the conservation of land will not be affected by the development. The Regulation also states that it is prohibited to straighten, change, divert or interfere in any way with the existing channel of a river, creek, stream or watercourse or change or interfere in any way with a wetland without permission from the Conservation Authority.

Per NVCA's interactive map viewer, a tributary of Beeton Creek enters the Subject Lands near the existing residence on the property. This tributary, and its associated floodplain, is identified as regulated area by the NVCA. Based on existing mapping, the regulated area is limited to the woodland in the northeastern corner of the property.

## 2.6 Provincial Endangered Species Act, 2007

The provincial ESA was developed to:

- Identify Species at Risk (SAR), based upon best available science;
- Protect SAR and their habitats and to promote the recovery of SAR; and
- Promote stewardship activities that would support those protection and recovery efforts.

The ESA protects all threatened, endangered and extirpated species listed on the Species at Risk in Ontario (SARO) list. These species are legally protected from harm or harassment and their associated habitats are legally protected from damage or destruction, as defined under the ESA.

# 3. Data Collection and Analyses

## 3.1 Background References

GEI has relied, in part, upon supporting background information to provide additional insight into the overall character of the Subject Lands. These resources included:

- MNRF Land Information Ontario (LIO) Natural Features Mapping;
- Natural Heritage Information Centre (NHIC) database;
- Provincial wildlife atlases (i.e., Ontario Breeding Bird Atlas, etc.);
- Citizen Science Databases (i.e., iNaturalist and eBird);
- DFO Aquatic Species at Risk Distribution Mapping; and
- Innisfil Creek Subwatershed Health Check 2023 (NVCA 2023).

The results of these background reviews are discussed in the following sections.

#### 3.1.1 Land Information Ontario

Based on the Ministry of Natural Resources and Forestry (MNRF) Land Information Ontario (LIO) geographic database, the following features were identified on or adjacent to the Subject Lands (**Figure 1, Appendix A**):

- Woodlands;
- Tributary of Beeton Creek; and
- ORMCP Natural Linkage Area.

## 3.1.2 Natural Heritage Information Centre

The Natural Heritage Information Centre (NHIC) database (MNRF 2023) was searched for records of provincially significant plants, vegetation communities and wildlife on, and in the vicinity of the Subject Lands. The database provides occurrence data by 1 km² area squares, with 4 squares overlapping at least a portion of the Subject Lands (17NJ9569, 17NJ9669, 17NJ9668, 17NJ9568). Within these squares, the search revealed 11 records, the following records are considered as current occurrences in this reporting:

- Species listed as Threatened or Endangered on the Species at Risk in Ontario (SARO) list:
  - o Eastern Meadowlark (Sturnella magna) Threatened in Ontario; and
  - o Bobolink (Dolichonyx oryzivorus) Threatened in Ontario;
- Species of Conservation Concern (i.e., listed as Special Concern on the SARO list, or identified as an S1-S3 species):
  - o Lilypad Clubtail (Arigomphus furcifer) S3 species
  - Nerveless Kuhlenberg's Sedge (Carex muehlenbergii var. enervis) S1/S2 species

#### 3.1.3 Ontario Breeding Bird Atlas

The Ontario Breeding Bird Atlas (OBBA) contains detailed information on the population and distribution status of Ontario birds (Bird Studies Canada et al. 2006). The data is presented on 100 km² area squares with one square overlapping a portion of the Subject Lands (17NJ96). It should be noted that the Subject Lands are a small component of the overall bird atlas square, and therefore it is unlikely that all bird species are found within the Subject Lands. Habitat type, availability and size are all contributing factors in bird species presence and use.

A total of 129 species were recorded in the atlas squares that overlap with the Subject Lands, with the following species of interest noted:

- Species listed as Threatened or Endangered on the SARO list:
  - o Bank Swallow (Riparia riparia) Threatened;
  - Bobolink Threatened;
  - Chimney Swift (Chaetura pelagica) Threatened;
  - Eastern Meadowlark Threatened;
  - Eastern Whip-poor-will (Antrostomus vociferus) Threatened;
  - Least Bittern (Ixobrychus exilis) Threatened; and
  - o Red-headed Woodpecker (*Melanerpes erythrocephalus*) Endangered.
- Species of Conservation Concern (i.e., listed as Special Concern on the SARO list, or identified as an S1-S3 species):
  - Barn Swallow (Hirundo rustica) Special Concern;
  - o Common Nighthawk (Chordeiles minor) Special Concern;
  - Eastern Wood-Pewee (Contopus virens) Special Concern;
  - o Grasshopper Sparrow (Ammodramus savannarum) Special Concern;
  - Golden-winged Warbler (Vermivora chrysoptera) Special Concern;
  - o Wood Thrush (*Hylocichla mustelina*) Special Concern;
  - o Common Gallinule (Gallinula galeata) S3B;
  - Purple Martin (Progne subis) S3B; and
  - Blue-winged Teal (Anas discors) S3B, S4M.

## 3.1.4 Ontario Reptile and Amphibian Atlas

The Ontario Reptile and Amphibian Atlas contains detailed information on the population and distribution status of Ontario herpetofauna (Ontario Nature 2020). The data are presented on 100 km<sup>2</sup> area squares with 1 square overlapping a portion of the Subject Lands (17NJ96). It should be noted that the Subject Lands are a small component of the overall atlas square, and therefore it is unlikely that all herpetofauna species are found within the Subject Lands. Habitat type, availability and size are all contributing factors in herpetofauna species presence and use.

A total of 20 species was recorded in the atlas squares that overlap with the Subject Lands, of which 4 are salamander species, 10 are frog and toad species, 2 are turtle species and 4 are snake species. Of these species, the following species of interest are noted:

- Species of Conservation Concern (i.e., listed as Special Concern on the SARO list, or identified as an S1-S3 species):
  - Snapping Turtle (Chelydra serpentina) Special Concern.

#### 3.1.5 Ontario Butterfly and Moth Atlas

The Ontario Butterfly and Moth Atlases (Toronto Entomologists' Association 2023, 2020), contain detailed information on the population and distribution status of Ontario butterflies and moths. The data are presented on 100 km² area squares with 1 square overlapping a portion of the Subject Lands (17NJ96). It should be noted that the Subject Lands are a small component of the overall atlas square, and therefore it is unlikely that all butterfly and moth species are found within the Subject Lands. Habitat type, availability and size are all contributing factors in butterfly and moth species presence and use.

A total of 63 species were recorded in the atlas squares that overlap with the Subject Lands, of which 44 are butterfly species and 19 are moth species. Of these species, only one species of conservation concern was identified; Monarch (*Danaus plexippus*), listed as Special Concern.

#### 3.1.6 Citizen Science Database (iNaturalist)

The iNaturalist (2023) database is a large citizen science-based identification and data collection app. It allows any citizen to submit observations to be reviewed and identified by other naturalists and scientists to help provide accurate species observations. As the observations can be submitted by anyone, and the records are not officially vetted, the data obtained from this tool should not be used as a clear indicator of species presence, and species may be filtered out based on habitat and target survey efforts.

This online database was examined to identify observations made within the Subject Lands that were research grade. However, no significant species were found on the Subject Lands or within 120 meters of its boundaries.

#### 3.1.7 Citizen Science Database (eBird)

The eBird (2023) database is a large citizen science-based project with a goal to gather bird diversity information in the form of checklists of birds, archive it, and share it to power new data-driven approaches to science, conservation and education. As the observations can be submitted by anyone, and the records are not officially vetted, the data obtained from this tool should not be used as a clear indicator of species presence, and species may be filtered out based on habitat and target survey efforts.

This online database was examined to identify observations made within the Subject Lands. However, no significant bird species were found on the Subject Lands or within 120 meters of its boundaries.

## 3.1.8 Aquatic Species at Risk Distribution Mapping

Aquatic species at risk distribution mapping (DFO 2023) was reviewed to identify any known occurrences of aquatic SAR, including fish and mussels, within the subwatershed where the Subject Lands are located.

No aquatic SAR habitats were identified on or within 120 m of the Subject Lands or within the subwatershed.

#### 3.1.9 Innisfil Creek Subwatershed Health Check 2023

The NVCA prepares watershed health checks, and a report was prepared for the Innisfil Creek subwatershed in 2023 (NVCA 2023). The report provides a high level overview of the state of the subwatershed, with the following observations noted:

- Forest cover within the subwatershed is generally poor, primarily due to the high quality of the land to support agricultural landscapes. Large stands are restricted to wetland communities (Cookstown Hollows Swamp and Bailey Creek Swamp Forest) and the scatted stands within the Oak Ridges Moraine, including those present on the Subject Lands. The woodland community that overlaps the southern portion of the Subject Lands is shown as containing interior forest habitat. The adjacent CUP woodland is also identified as 'forest gained' between 2008 and 2018. In addition, a natural corridor for wildlife movement is shown generally in an east-west direction through the woodland communities across the Oak Ridges Moraine, with the greatest connectivity shown north of the Subject Lands.
- Wetland habitats within the subwatershed are generally fair to poor when compared to Environment Canada's targets; no wetlands are identified in the vicinity of the Subject Lands.
- The Subject Lands eventually drain through tributaries towards Beeton Creek near Tottenham to the north of the Subject Lands. Beeton Creek is generally mapped as impaired to below potential stream health.
- The Subject Lands are mapped as being within a significant groundwater recharge area

#### 3.2 Technical Methods and Field Studies

A site reconnaissance visit was completed in June 2021 to assess ecological conditions on the Subject Lands. A scoped ecological field survey program was undertaken in 2022. The following ecological surveys have been completed within the Subject Lands:

- Site Reconnaissance;
- Three-season Botanical Inventories and Ecological Land Classification (ELC);
- Headwater Drainage Feature Assessment;
- Breeding Bird Surveys; and
- Incidental Wildlife Observations (recording during other surveys).

A summary of the dates and weather condition associated with all the aforementioned surveys is provided in **Table 1** (**Appendix B**).

Both the NHIC (MNRF 2022) database and the Species at Risk in Ontario (SARO) list (O.Reg. 230/08) were reviewed to determine the current provincial status for all species identified during the field programs outlined below.

## 3.2.1 Vegetation and ELC Methods

ELC surveys were completed in June 2021, May 2022, July 2022 and September 2022. Vegetation community types were initially assessed through the use of aerial imagery, and were confirmed, sampled and revised, as necessary, using the sampling protocol of the

ELC Southern Ontario (Lee et al., 1998). ELC was completed to the finest level of resolution (Vegetation Type) where feasible. Species names generally follow nomenclature from the Database of Vascular Plants of Canada (Brouillet et al., 2010+).

The provincial status of all vegetation communities is based on NHIC (MNRF 2022). Identification of potentially sensitive native plant species is based on their assigned coefficient of conservatism (CC) value, as determined by Oldham et al. (1995). The CC value, ranging from 0 (low) to 10 (high), is based on a species tolerance of disturbance and fidelity to a specific natural habitat. Species with a CC value of 9 or 10 generally exhibit a high degree of fidelity to a narrow range of habitat parameters.

## 3.2.2 Headwater Drainage Feature Assessment

Potential headwater drainage features (HDFs) on the Subject Lands were assessed using the TRCA/CVC (2014) "Evaluation, Classification and Management of Headwater Drainage Features Guidelines" (herein referred to as the HDFA Guidelines). These guidelines provide a standardized means of identifying and assessing the value of HDFs and identifying long-term management recommendations to protect or maintain the important ecological or biophysical functions provided by HDFs in a developing landscape.

Per the requirements of the HDFA Guidelines, GEI completed site visits at seasonally appropriate times to verify the presence of any features identified through a review of aerial imagery, provide up-to-date data regarding existing HDF conditions and ensure that a full data set is available. HDFA surveys are completed in accordance with the protocols in the HDFA Guidelines, with up to 3 site visits potentially required.

During the first site visit, all areas of the Subject Lands were walked to identify potential HDFs. Consideration was particularly given to areas identified as containing low ground based on topographic information. No HDFs were identified on the Subject Lands.

However, the mapped tributary to Beeton Creek was also inspected during this initial visit, as well as a subsequent site inspection completed in late May 2022 following a precipitation event. It was determined to meet the criteria of a drainage feature and not a watercourse. Further details are provided in **Section 4.5**.

## 3.2.3 Breeding Bird Surveys

Breeding bird surveys were conducted following protocol set forth by the Ontario Breeding Bird Atlas (Cadman et al. 2007) and the Ontario Forest Bird Monitoring Program (Cadman et al. 1998).

Surveys were conducted between dawn and five hours after dawn with suitable wind conditions, no thick fog or precipitation (Cadman et al., 2007). Point count stations were located in various habitat types within the Subject Lands and combined with area searches to help determine the presence, variety and abundance of bird species. Each point count station was surveyed for 10 minutes for birds within 100 m and outside 100 m. All species recorded on a point-count were mapped to provide specific spatial information and observed for signs of breeding behaviour.

# 4. Bio-Physical Characterization

## 4.1 Physiography

The Subject Lands are located on the Oak Ridges Moraine (ORM), an important geological landform that was formed through the advancement and retreat of glaciers, and the deposition of stratified sediment. The ORM and underlying sediments are late Pleistocene in age and overlie thin Paleozoic bedrock platform strata.

The Subject Lands reach a high point at the woodland communities at the southern extent of the Subject Lands, and generally the land falls away to the north, west and east of that point, with some rolling topography present across the Subject Lands.

## 4.2 Landscape Ecology

Situated within the Oak Ridges Moraine, movement predominantly occurs through the landscape in an east-west direction through the various woodlands. Larger communities in relative proximity to the Subject Lands are predominantly to the west, with movement occurring towards Palgrave and the Palgrave Forest and Wildlife Area to the north and Albion Hills Conservation Park to the south of the residential developments in that area.

Movement to the south through the scattered woodlands eventually connects to the Humber River and the surrounding valleylands that would connect to the south towards Lake Ontario.

## 4.3 Vegetation

## 4.3.1 Ecological Land Classification

The Subject Lands were dominated by tilled agricultural lands, that were planted with soybean in 2022. Surrounding these agricultural lands, two blocks of mature woodland were present. These features were classified Fresh-Moist Sugar Maple – Hardwood Deciduous Forest (FOD6-5). These woodlands also extend beyond the Subject Lands to the northeast and south. It is noted that the connection between these two woodlands have been fragmented.

The canopy of these mature forests was dominated by Sugar Maple (*Acer saccharum*), with some Eastern Hemlock (*Tsuga canadensis*) and a small number of Eastern White Pine (*Pinus strobus*). The sub-canopy was dominated by Sugar Maple with Green Ash (*Fraxinus pennsylvanica*) and American Beech (*Fagus grandifolia*). The shrub layer was similarly dominated by Sugar Maple, Green Ash, though Red Elderberry (*Sambucus racemosa*) was also abundant. The ground layer was diverse, some dominant species noted included: Virginia Waterleaf (*Hydrophyllum virginianum*), Canada Wild-Ginger (*Asarum canadense*), Wild Leek (*Allium tricoccum var. tricoccum*), Canada Enchanter's Nightshade (*Circaea canadensis ssp. canadensis*), Garlic Mustard (*Alliaria petiolata*) and Herb-Robert (*Geranium robertianum*). Where moister soil conditions were available, Ostrich Fern (*Matteuccia struthiopteris var. pensylvanica*) and Spotted Jewelweed (*Impatiens capensis*) were more dominant.

In addition to the woodlands, two small hedgerows were also present within the Subject Lands. The south-most hedgerow consisted mainly of Sugar Maple, American Basswood (*Tilia americana*) as well as Ash (*Fraxinus spp.*) snags. While the north-most hedgerow mainly consisted of Scots Pine (Pinus sylvestris) with Common Buckthorn (*Rhamnus catharica*), Tatarian Honeysuckle (*Lonicera tatarica*).

Immediately west of the Subject Lands, a small Coniferous Cultural Plantation (CUP) was present. This feature was situated centrally to the Subject Lands. To the north-northwest of this feature, a Mineral Cultural Meadow (CUM1) was present that extended to Hunsden Sideroad.

These vegetation communities are shown on Figure 2 (Appendix A).

#### 4.3.2 Vascular Plants

Botanical inventories completed on the Subject Lands have identified a total of 152 species of vascular plants. Of that number, 103 (or 68%) were native and 49 (or 32%) are exotic. A completed species list is provided in **Table 2** (**Appendix B**).

The majority of the native species (98%) were ranked S5 (secure in Ontario) or S4 (apparently secure in Ontario; NHIC, 2017). One species is listed as an S2? (rare in Ontario); Butternut (*Juglans cinerea*), which are also listed as Endangered on the SARO List. A total of four Butternut were identified within the Subject Lands, these were all associated with Fresh-Moist Sugar Maple – Hardwood Deciduous Forest (FOD6-5) located in the northeast of the Subject Lands (**Figure 2**, **Appendix A**).

#### **Locally Rare Plants**

In addition to Butternut, 12 regionally rare or uncommon plants were observed, as per the Peel Region, rarity rankings (Varga 2005). None of these regionally rare species are considered rare in Ontario. None of the species recorded from the Subject Lands had a coefficient of conservation value of 9 or 10. The regionally rare species are summarized below:

- Rare Species:
  - Downy Arrowwood (Viburnum rafinesquianum);
  - Blue Cohosh (Caulophyllum thalictroides);
  - Common Bedstraw (Galium aparine);
  - White Bear Sedge (Carex albursina);
  - Loose-Flowered Sedge (Carex leptalea);
  - Sprengel's Sedge (Carex sprengelii); and
  - Interrupted Fern (Osmunda claytoniana).
- Uncommon Species:
  - Virginia Stickseed (Hackelia virginiana);
  - Squirrel-Corn (Dicentra canadensis);
  - o Canada Plum (*Prunus nigra*);
  - o Rough Bedstraw (Galium asprellum); and
  - Bristle-Stalked Sedge (Carex leptalea).

These species were identified within forest communities on the Subject Lands.

#### **Invasive Plants**

Invasive plants are those that can become (or presently are) a serious problem within a defined location. These plants reproduce and spread aggressively, reducing the local biodiversity and threatening ecological function. Depending on existing conditions, some invasive species can outcompete all other species.

Urban Forest Associates (2002) provides a categorical ranking system for plants known to be invasive in southern Ontario. Of the 49 exotic species observed on the Subject Lands, six are ranked as Category 1 by Urban Forest Associates.

Category 1 plants are deemed to be the most invasive and can dominate a site indefinitely. These are a threat to natural areas wherever they occur because they have very effective reproduction and dispersal mechanisms. The Category 1 plants observed on the Subject Lands are:

- European Swallowwort (Vincetoxicum rossicum);
- Garlic Mustard:
- Dame's Rocket (Hesperis matronalis):
- Tartarian Honeysuckle (Lonicera tatarica);
- European Buckthorn (Rhamnus cathartica); and
- Manitoba Maple (Acer negundo).

#### 4.4 Wildlife

## 4.4.1 Breeding Bird Surveys

A total of 39 bird species were observed within the Subject Lands. Of this total, two species are confirmed, 21 are probable, and 13 are possible breeders on the Subject Lands. The remaining three bird species are considered non-breeders, flyovers, or migrants. An additional twelve species were observed only on surrounding lands within 120 m. The observed breeding bird species are discussed in the sections below. All species observed on the Subject Lands are listed in **Table 3** (**Appendix B**).

All (100%) of the confirmed, probable or possible breeders are provincially ranked S5 (common and secure), S4 (apparently common and secure) or SNA (species not native to Ontario). No bird species are considered provincially rare (S1-S3; NHIC 2021) However, the following Species of Conservation Concern were observed on the Subject Lands associated with the woodland communities:

- Wood Thrush Special Concern; and
- Eastern Wood-Pewee Special Concern.

Additionally, the following Species at Risk and Species of Conservation Concern were observed beyond the Subject Lands within suitable meadow habitats, however it should be noted that the Subject Lands do not provide breeding habitat for these species:

- Eastern Meadowlark Threatened; and
- Grasshopper Sparrow Special Concern.

#### 4.4.2 Incidental Wildlife

A total of 20 species were recorded incidentally within the Subject Lands. Of these, one SAR bird, a Bobolink, was recorded within the Subject Lands. Bobolink is designated as Threatened in Ontario, the remaining species observed are considered common or non-native to Ontario.

The observation of Bobolink was made on June 2, 2021, during the site reconnaissance visit, and was observed alighting on a small tree. Given that the Subject Lands are comprised of soy crops and forested habitat, suitable nesting habitat for the species was not identified within the Subject Lands, and this species was not recorded during 2022 surveys.

A summary of all wildlife observed incidentally is provide in **Table 4** (**Appendix B**).

## 4.5 Headwater Drainage Features

A site visit was completed on April 20, 2022, to assess for presence of drainage features on the Subject Lands. Agricultural fields had been recently tilled prior to the site investigation; however, no evidence of water movement or presence was observed on the site.

A mapped tributary to Beeton Creek is shown on NVCA mapping within the northeastern extent of the Subject Lands, near the existing residence. In conjunction with HDFA Round 1, the mapped tributary was also inspected. The arterial branch to the south was determined not to exist on the landscape. The main branch of the feature was characterized as a poorly defined drainage swale with stagnant and isolated pools of water located towards the upstream end of the Subject Lands. The feature connects beneath Hunsden Road through a corrugated steel pipe (CSP) which was also dry during the time of the initial inspection. A subsequent site inspection was completed following a precipitation event in late May 2022 and further confirmed the drainage feature to be dry. Given the lack of flow and very poor channel definition associated with the drainage swale, the feature is proposed to be evaluated and managed for under the HDFA Guidelines. This determination is also supported by the fluvial geomorphological assessment which determined that this feature should not be identified as a watercourse (GEI 2023). Classification of the feature is provided in the table below.

STEP 1. HYDROLOGY		STEP 2. RIPARIAN	STEP 3. FISH HABITAT	STEP 4. TERRESTRIAL HABITAT	MANAGEMENT RECOMMENDATION
FUNCTION	MODIFIERS				
FT – 7 FC – 2 (Round 1) FC – 1 (Round 2)  Recharge Functions (based on location in groundwater recharge area) – dry or standing water was identified during assessments.		Important – forest habitat adjacent to feature.	Contributing- No direct fish habitat is present.	Limited – As per Table 7 in HDFA Guidelines, swale provides limited terrestrial function Although this feature is situated within a forest, it does not connect other features upstream and downstream.	Maintain Recharge

Evaluation of the NVCA mapped tributary under the HDFA guidelines results in a management recommendation of *Maintain Recharge*. This management recommendation includes the following strategy:

• Maintain overall water balance by providing mitigation measures to infiltrate clean stormwater.

As the feature is well protected within the woodland, there are no material ecological requirements with respect to this feature. Water balance measures at the site level will ensure recharge functions are maintained.

# 5. Analysis Of Ecological and Natural Heritage Significance

Eight types of natural features are identified in the PPS (MMAH 2020):

- Significant wetlands;
- Significant coastal wetlands;
- Significant woodlands;
- Significant valleylands;
- SWH;
- Fish habitat;
- · Habitat of endangered and threatened species; and
- Significant areas of natural and scientific interest.

In addition to natural features identified in the PPS, consideration was also given to the KNHF and KHF identified within the ORMCP, as well as woodlands that may meet criteria to be identified as Core Areas, Natural Areas and Corridors or Potential Natural Areas and Corridors under the Region of Peel Official Plan. Finally, an assessment with respect to which areas meet the criteria for identification as EZ1 or EZ2 was completed.

The presence/absence of these natural features on the Subject Lands are discussed in the subsequent sections of this EIS. The NHRM (MNR 2010) and the Peel-Caledon Significant Woodlands and Significant Wildlife Habitat Study (NSE Inc et al 2009) were referenced to assess the potential significance of these feature types.

Where natural features are present on the Subject Lands, their sensitivities are discussed.

## 5.1 Wetlands/Significant Wetlands

Within Ontario, Significant Wetlands are identified by the MNRF or by their designates. Other evaluated or unevaluated wetlands may be identified for conservation by the municipality or the conservation authority.

No significant wetlands are known to occur within 750 m of the Subject Lands, with the nearest provincially significant wetland (PSW), known as the Gibson Lake Wetland Complex, identified more than 1 km to the West. No wetlands were identified on the Subject Lands during the site investigations. As a result, this feature type is not present.

## 5.2 Significant Coastal Wetlands

Significant coastal wetlands are not present on or adjacent to the Subject Lands.

## 5.3 Habitat of Endangered and Threatened Species

Based upon the species identified through the background wildlife atlas search (**Section 3.0**) and field investigations results, the following SAR and SAR habitat is confirmed or may be present within the Subject Lands:

- Butternut;
- Eastern Small-footed Myotis (Myotis leibii);
- Little Brown Myotis (Myotis lucifugus);
- Northern Myotis (Myotis septentrionalis); and
- Tricolored Bat (Perimyotis subflavus).

Of these, only Butternut was confirmed within the Subject Lands, associated within the northeastern woodland community. In addition, the forested habitats within the Subject Lands are considered candidate habitat for Eastern Small-footed Myotis, Little Brown Myotis, Northern Myotis, and Tricolored Bat, and are thus treated as habitat for Endangered bat species. Details pertaining to these species are provided below. The remaining species identified from the records review were confirmed absent, or the Subject Lands did not provide suitable habitat for the species.

Both Bobolink and Eastern Meadowlark, two SAR bird species that breed in grassland habitats, were recorded incidentally during surveys on or surrounding the Subject Lands. However, as breeding habitat for these species is not present within the Subject Lands, there will be no impacts to the habitat for these species, and they are not addressed further.

#### 5.3.1 Butternut

Four Butternut were identified within the northeastern woodland community on the Subject Lands. Butternut are listed as Endangered on the SARO List. Of the four trees, two of the trees were identified along the southern extent of the forest edge and two were set further back into the feature. The two trees located near the forest edge are more than 30 m from the proposed structural envelope (**Figure 2**, **Appendix A**), and abut active agricultural fields.

#### 5.3.2 Endangered Species of Bats

Suitable bat maternity roosting habitat (tree cavities, peeling bark, rock piles etc.) are present within the woodland communities on the Subject Lands. Targeted surveys for bats were not completed as no encroachment into these features are proposed. As a result, the woodlands are treated as candidate SAR bat habitat.

#### 5.4 Fish Habitat

Fish habitat, as defined in the federal Fisheries Act, c. F-14, means... spawning grounds and nursery, rearing, food supply, and migration areas on which fish depend directly or indirectly to carry out their life processes. Fish, as defined in S.2 of the Fisheries Act, c. F-14, includes parts of fish, shellfish, crustaceans, marine animals and any parts of shellfish, crustaceans or marine animals, and the eggs, sperm, spawn, larvae, spat and juvenile stages of fish, shellfish, crustaceans and marine animals.



Though no water movement was observed during site investigations completed in 2022, the drainage feature present in the northeastern portion of the Subject Lands may be considered to provide indirect fish habitat. Provided that the conveyance of any potential seasonal or event based flows are still directed towards the culvert beneath Hunsden Sideroad, no negative impacts to potential fish habitat are anticipated.

## 5.5 Significant Woodlands/Region of Peel Woodlands

The PPS notes that, significant woodlands should be defined and designated by the planning authority using criteria established by the MNRF. Within the Region of Peel, woodlands are assessed against criteria to determine whether they are considered to be Core Areas of the Greenlands System, Natural Areas and Corridors (NAC) of the Greenlands System, or Potential NAC (PNAC) using criteria identified in Table 1 of the Official Plan. As the Subject Lands are situated within the Oak Ridges Moraine Conservation Plan Area, consideration was also given to the criteria identified in Technical Paper 7 of the ORMCP.

Both woodland communities on the Subject Lands were assessed against these criteria.

The southern woodland is a large feature that is well more than 16 ha in size. Based on size alone, this feature would be considered a Core Area under both the ROP and Technical Paper 7, and so would be considered a significant woodland.

The northeastern woodland on the Subject Lands is approximately 6 ha in size. As this woodland community contains Butternut, a species listed as Endangered by both the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) and the Committee on the Status of Species at Risk in Ontario (COSSARO), this feature would also be considered to be a Core Area under the ROP, and also meets criteria for significance under Technical Paper 7.

The limits of these features were staked with the Town of Caledon and members of the Project Team. It is noted that the staking included portions which are not considered to be components of the significant woodland. This included a small hedgerow extending southwesterly from the northeastern woodland, along the limit of the previously severed residential lot along Hunsden Sideroad, as well as a single mature tree in the northwest corner of the southern woodland that is a portion of the remnant hedgerow that extends further to the northwest. These features are not considered to be components of the significant woodland and setbacks from these features are not provided.

The cultural plantation west of the Subject Lands was also considered. The feature measures approximately 1.5 ha in size, with several large openings within the canopy. As this feature is not identified within either the ORMCP Natural Core or Natural Linkage Areas (see discussions re linkage areas in section 5.13) and is not situated within nor intersects with a key natural heritage feature or hydrologically sensitive feature or their vegetation protection zone, this woodland is neither a core area nor a significant woodland under the ORMCP. However, as the cultural plantation is greater than 1.5 ha in size, it does exceed the size requirement to be identified as a PNAC.

## 5.6 Significant Valleylands

Significant Valleylands should be defined and designated by the planning authority. General guidelines for determining significance of these features are presented in the NHRM) for Policy 2.1 of the PPS. Recommended criteria for designating significant valley lands include prominence as a distinctive landform, degree of naturalness, and importance of its ecological functions, restoration potential, and historical and cultural values.

No valley features were identified on the Subject Lands.

## 5.7 Significant Areas of Natural and Scientific Interest (ANSIs)

An ANSI is identified by the MNRF as "areas of land and water containing natural landscapes or features that have been identified as having life science or earth science values related to protection, scientific study or education" (MNR 2010).

A review of mapping from MNRF's LIO and NHIC databases showed that there are no ANSIs identified on or in proximity to the Subject Lands.

## 5.8 Significant Wildlife Habitat

Significant wildlife habitat (SWH) is one of the more complex natural heritage features to identify and evaluate. There are several provincial documents that discuss identifying and evaluating SWH: the NHRM, the SWH Technical Guide (MNR 2000), and the relevant SWH Ecoregion Criterion Schedule (MNRF 2015). The Subject Lands are located in Ecoregion 6E and were therefore assessed using the 6E Criterion Schedule (MNRF 2015).

There are four general types of significant wildlife habitat, including seasonal concentration areas of animals, migration corridors, rare vegetation communities or specialized habitat for wildlife, habitat for species of conservation concern and animal movement corridors. All types of significant wildlife habitat are discussed in more detail below.

#### Seasonal Concentration Areas of Animals

Seasonal concentration areas are those sites where large numbers of a species gather together at one time of the year, or where several species congregate. The following is a partial list of numerous examples: deer yards, snake and bat hibernacula, waterfowl staging areas, raptor wintering areas, bird nesting colonies, shorebird stopover areas, and colonial nesting bird habitats. Areas that support a SAR, or if a large proportion of the population may be lost if the habitat is destroyed, are examples of seasonal concentration areas which should be designated as significant.

#### Rare Vegetation Communities or Specialized Habitat for Wildlife

Rare vegetation communities or specialized habitat are two separate components.

Rare vegetation communities are those that are considered rare in the province. These include cliff and talus slopes, sand barrens, alvars, old growth forest, savannah, and tallgrass prairie. Provincially ranked vegetation communities with SRANKS of S1 to S3 (extremely rare to rare-uncommon in Ontario) as defined by the NHIC, would also typically qualify. It is assumed that these habitats are at risk and that they are also likely to support additional wildlife species that are considered significant. Such vegetation communities do not occur on or adjacent to the Subject Lands.

Specialized habitats are microhabitats that are critical to some wildlife species. The NHRM defines specialized habitats as those that provide for species with highly specific habitat requirements; areas with exceptionally high species diversity or community diversity; and areas that provide habitat that greatly enhances species' survival.

#### Habitat for Species of Conservation Concern

Habitat for species of conservation concern includes five types of habitats:

- a) Marsh bird breeding habitat;
- b) Open country bird breeding habitat;
- c) Shrub/early successional bird breeding habitat;
- d) Terrestrial crayfish; and
- e) Special concern and rare wildlife species.

Habitats of species of conservation concern do not include habitats of Endangered or Threatened species as identified by the Endangered Species Act, 2007, which are discussed in Section 5.3.

#### **Animal Movement Corridors**

Animal movement corridors are areas that are traditionally used by wildlife to move from one habitat to another. This is usually in response to different seasonal habitat requirements. Some examples are trails used by deer to move to wintering areas, and areas used by amphibians between breeding and summering habitat. Animal movement corridors are only identified as significant wildlife habitat where a confirmed or candidate significant wildlife habitat has been identified by MNRF or the planning authority.

**Table 5** (**Appendix B**) discusses the potential for SWH within the Subject Lands based on the background information review and field investigations. The following SWH types were identified within the Subject Lands:

- Candidate Bat Maternity Colonies (FOD6-5);
- Candidate Reptile Hibernacula (rock pile within FOD6-5);
- Confirmed Woodland Area-Sensitive Bird Breeding Habitat (southernmost FOD6-5);
- Confirmed Habitat for Species of Conservation Concern:
  - Eastern Wood-Pewee; and
  - Wood Thrush.



## 5.9 Sand Barrens, Savannahs and Tallgrass Prairies

These vegetation types were not identified on the Subject Lands.

#### 5.10 Permanent and Intermittent Streams

As discussed in section 4.5, there is no permanent or intermittent stream present on the Subject Lands.

#### 5.11 Kettle Lakes

Kettle lakes are not present on the Subject Lands.

## 5.12 Seepage Areas and Springs

No seepage areas or springs were identified on the Subject Lands.

## 5.13 Linkage Assessment

The Oak Ridges Moraine Conservation Plan identifies Natural Linkage Areas on the Subject Lands, consisting of an east-west corridor along Hunsden Sideroad, and a north south corridor connecting the two woodlands that overlap the Subject Lands. This Natural Linkage Area was later refined in accordance with the provisions of the ORMCP within the Town of Caledon Official Plan to reduce the extent of the Natural Linkage Area on the Subject Lands. Schedule G of the Official Plan shows the east-west linkage corridor predominantly along the northern side of Hunsden Sideroad (as delineated by the extent of Policy Area 4 on Schedule G, which delineates the extent of ORMCP Natural Core and Linkage Areas). Given the increased prevalence of trees/shrubs along the northern side of Hunsden Sideroad, this is the most probable location for wildlife movement across the landscape. Along the southern side of Hunsden Sideroad, there is a general absence of vegetation cover, with several large gaps between trees. Wildlife movement along the southern side of Hunsden Sideroad is less preferred when compared to the northern side. Preservation of a linkage along the southern side of Hunsden Sideroad would not be warranted; as a result, the recommended location for the linkage remains on the north side of Hunsden Sideroad, consistent with the mapping shown in Schedule G of the Town of Caledon Official Plan.

The other location shown for a linkage is along the eastern property limit between the northern and southern woodland communities. A projection of the northern woodland extends towards the southern woodland along the adjacent property to the east. A 60 m gap exists between the two woodland communities. The natural linkage area within this area would exist through the woodland communities, and along the narrow remnant vegetation along the property line within the gap between the two features.

## 5.14 EZ Designations

As identified within **Section 2.1**, the following are identified as the requirements with respect to EZ designations:

- EZ1 designations includes all ORMCP KNHF and KHF, and their related Minimum Vegetation Protection Zones (MVPZ), along with more sensitive biological communities, valley and stream corridors and their associated floodplains, native upland and lowland woodlands, natural waterbodies, Provincially and locally significant wetlands, and Environmentally Significant/Sensitive Areas along with other features of local or regional importance.
- EZ2 Designations are locations with high groundwater table, seasonal flooding, dry swale lowlands and natural depressions performing natural run-off, detention and groundwater recharge functions, and smaller hedgerows and strips of native vegetation.

Within the Subjetc Lands, as identified in Schedule I of the TCOP (2018), the significant woodlands (FOD6-5) and the Cultural Plantation (CUP) are designated as EZ1s.

The Town of Caledon was consulted to determine what information resulted in the identification of an EZ2 on the Subject Lands, however no information is currently available. As a result, an assessment was completed to determine whether that portion of the Subject Lands meets the definition of an EZ2, as discussed further below.

- High groundwater table: High groundwater tables are defined in the Town of Caledon Official Plan as areas where the water table is usually within 1.5 m or less of the ground surface. The groundwater table on the site is greater than 1.5 m below the ground surface.
- Seasonal flooding: Investigations in the spring did not find any evidence of seasonal flooding on the Subject Lands.
- Dry swale lowlands and natural depressions performing natural run-off detention and groundwater recharge functions: A review of the topographic mapping showed that there are no topographic lowlands present in this area that would provide natural run-off detention and groundwater recharge functions.
- Smaller hedgerows and strips of native vegetation: There are no smaller hedgerows
  or strips of native vegetation present in the area of EZ2 in the center of the Subject
  Lands, however an EZ2 is likely present associated with the hedgerow connecting the
  forest units along the eastern property limit. This area is generally already contained
  within the VPZ of the larger woodlands and was discussed above with respect to the
  linkage assessment.

Given the above assessment, it is the recommendation of this NHE that there is no EZ2 designation present within the central portion of the Subject Lands, and as a result, the restrictions associated with that designation would not apply to the proposed development.

# 5.15 Summary of Ecological Components Subject to Impact Assessment

An analysis of existing natural heritage features on the Subject Lands was completed. The results of this analysis identified the following natural heritage features as present, within and/or adjacent to the Subject Lands:

- Habitat of Endangered and Threatened Species (Butternut, Bat SAR), associated with the deciduous forest communities on and adjacent to the Subject Lands;
- Significant Woodlands/Core Area Woodlands/EZ1 area associated with the deciduous forest communities on and adjacent to the Subject Lands;
- PNAC Woodland/EZ1 area associated with the cultural plantation west of the Subject Lands:
- Indirect Fish Habitat associated with the drainage feature on the Subject Lands;
- Candidate and Confirmed SWH associated with the deciduous woodland communities on the Subject Lands:
  - Candidate Bat Maternity Colonies (both FODs);
  - o Candidate Reptile Hibernacula (south-most FOD6-5);
  - Confirmed Woodland Area-Sensitive Bird Breeding Habitat (south-most FOD6-5);
  - o Confirmed Habitat for Species of Conservation Concern (both FODs):
    - Eastern Wood-Pewee; and
    - Wood Thrush.
- Linkage corridors along the northern extent of Hunsden Sideroad and along the eastern portion of the Subject Lands between the two significant woodlands.

These features are shown on Figure 3 (Appendix A).

# 6. Description Of Proposed Development

The proposed development would consist of estate residential development within the existing agricultural lands.

The Draft Plan (Appendix C) considers 13 estate residential lots of approximately 1.2 to 1.8 acre in size, connected via a Street "B" connection to the development presently in construction to the west, and a Street A connection to Hunsden Sideroad. Street "A" is presently proposed to terminate at a cul-de-sac within the southeastern extent of the Subject Lands, though an open space corridor is maintained that would permit an eventual connection to the lands to the east if required. The Draft Plan proposes a 30 m setback from the woodland communities on the Subject Lands, consisting of a 20 m vegetated buffer outside of the proposed lots, along with a 10 m setback for the structural envelope within the lots.

A Functional Servicing and Preliminary Stormwater Management Report was prepared by C.F. Crozier & Associates Inc. (dated November 2023). Within the report, it has been recommended that roadside bioswales and a rear yard infiltration trench will be used to control peak flows post-development to the pre-development levels up to the 100-year storm events. In addition, the bioswales will incorporate filter media and grassed swales to control water quality.

# 7. Impact Assessment, Mitigation and Enhancement Opportunities

This section of the report assesses the potential effects on the natural heritage features and their associated functions that could occur over the short-term and long-term following implementation of a future development plan, based on the draft concept plan. It also suggests appropriate mitigation measures to limit negative impacts and/or to enhance features and functions where practical.

The impact assessment below examines the predicted effects of the development on the natural heritage features and associated functions present on, and adjacent to, the Subject Lands. (**Figure 4, Appendix A**) shows the proposed development against the identified features.

## 7.1 Vegetation Protection Zone

This section provides a technical summary regarding the recommended vegetation protection zone for the identified natural heritage features on the Subject Lands. These are discussed by feature type.

#### 7.1.1 Significant Woodlands

Two significant woodland units are present on the Subject Lands. The significant woodlands were also assessed as being habitat for endangered species (Butternut, endangered species of bats), as well as both confirmed and candidate significant wildlife habitats.

The key functions of these areas on the Subject Lands are:

- Provision of breeding, foraging, and wintering habitat for wildlife, including birds, mammals, reptiles, and amphibians, including endangered species of bats and special concern wildlife species;
- Provision of habitat for the endangered Butternut;
- Provision of movement corridors for wildlife with connections to other natural core areas within the Oak Ridges Moraine area;
- Nutrient cycling and hydrological cycling associated with woodland vegetation communities; and
- Improvements in air quality relating to the presence of woodland vegetation.

Understanding those features and functions, allows for the consideration of an appropriate protective buffer. The recommended buffers or vegetation protection zones, has considered the physical characteristics of the natural edge (e.g., soil types, moisture regime, woodland edge structure and composition and rooting habits of adjacent trees) and has addressed other pertinent aspects.

In accordance with relevant policies, a 30 m development setback has been incorporated into the development plan, consisting of a 20 m wide buffer block (Block 16) that will be subject to reforestation plantings (discussed further in section 8) and a 10 m wide portion within each of the proposed lots abutting this feature situated outside of the structural envelope.

This development setback has been assessed and is considered appropriate as outlined below:

- Given the use of the adjacent agricultural fields that are proposed for development as row crop agricultural, these lands do not provide critical support functions for wildlife within the significant woodlands, which would necessitate the maintenance of larger buffers for protection, as noted below:
  - General wildlife use of the lands adjacent to the significant woodlands is likely restricted to periodic transit through, or foraging opportunities within, given the existing conditions.
  - o Bat species may forage over the agricultural lands; however, alternate foraging habitat is abundant in the local area. In addition, with the estate residential development, open landscaped areas will remain on the landscape that would also provide foraging opportunities. The SWH Mitigation Support Tool (MNRF 2014), does not recommend a specific setback for maternity colony habitats.
  - Reptile movement from the candidate hibernacula feature will continue throughout the landscape. The SWH Mitigation Support Tool (MNRF 2014), states that, the SWH includes a 30 m radius around the hibernacula feature. This radius is protected within the woodland buffers. Further, there is no development proposed within the woodland communities which would provide necessary shelter cover for aggregating snakes prior to entry into the hibernacula.
  - There will be no impact to the structure of the woodland communities providing area-sensitive woodland bird breeding habitat, and habitat for Wood Thrush and Eastern Wood-Pewee. The SWH Mitigation Support Tool (MNRF 2014), does not recommend a specific setback for these habitat types. The measures to protect the significant woodland will be effective at protecting this habitat type.
- There would be no direct impact to the Butternut trees required within the woodland, and as the fields are in agricultural production, there is no reproductive dispersal habitat available in the present condition beyond the limits of the woodland.
- Mitigation measures with respect to soil erosion, stormwater management, and water balance are addressed separately in the sections below.
- No vegetation removal is proposed, and therefore there would be no anticipated impact on nutrient cycling or air quality benefits within this feature.

#### 7.1.2 PNAC Woodland

A PNAC Woodland was identified to the west of the Subject Lands. This woodland is a cultural plantation and has not been identified as being associated with any other natural heritage features. Given their monocultural planting regime, cultural plantations such as this one are generally of low ecological function. Based on the site investigations, this is considered applicable to the feature in question.

Given the above, a 10 m development setback is proposed from the staked limits of this feature. The development setback will be located within each of the proposed lots abutting

this feature situated outside of the structural envelope. The structural setback has been aligned to be 15 m from the property limit, which results in a variable buffer greater than 10 m of approximately 12 to 15 m depending on the extent of the overhang of the vegetation along the property limit. The area between the property limit and the structural setback will be seeded with a native seed mix

#### 7.1.3 Linkages

There are two linkages identified within proximity of the Subject Lands.

The first feature is located north of Hunsden Sideroad. As the feature is on the other side of the road, a development setback is not recommended for this feature.

The second feature was identified encompassing the two significant woodlands, with a connection between these two features. The only existing connection between these woodlands at present is a small linear connection along the property line with marginal vegetation. To protect and enhance/restore this connection, a 20 m vegetation protection zone has been proposed between the two significant woodlands as a movement corridor between these features. Given the open nature of this connection, this 20 m vegetation protection zone is proposed to be subject to reforestation measures as outlined in section 8, which will represent a substantial improvement to connectivity between these features.

As discussed above, an open space corridor has been provided that would permit future construction of a roadway though this area to the adjacent lands to the east. Given both the proximity and general narrowness of the connection between these two features, this section of roadway would likely see significantly higher wildlife movement than other connections on the landscape. If a roadway is placed through this area, the design should consider the potential for increased wildlife/vehicle interactions, the need for provision of wildlife passages beneath the roadway should be assessed, and appropriate signage displayed.

#### 7.2 Indirect Fish Habitat

#### 7.2.1 Potential Impacts During Construction

As the drainage feature is situated within the woodland community away from the proposed development, no direct impacts would occur.

Potential indirect impacts could occur during construction include:

- Erosion and sedimentation from the construction area;
- Effects due to stormwater management during construction; and
- Accidental spills (e.g., fuel or oil from machinery) with transport of spilled material to watercourses.

Each of these potential impacts is discussed in the following sections.

#### **Erosion and Sedimentation**

Erosion and sedimentation from the disturbed work area associated with the proposed development could potentially result in adverse effects to water quality (e.g., increased turbidity) or sedimentation and associated effects on fish (e.g., injury or mortality due to suspended sediments or altered habitat use) or fish habitat (e.g., loss of interstitial spaces in

rocky areas, smothering of aquatic vegetation and/or incubating eggs) in downstream receiving features.

A preliminary Erosion and Sedimentation Control (ESC) Plan has been prepared by C.F. Crozier & Associates to minimize the potential for erosion and sedimentation from the construction site. The preliminary plan includes the use of sedimentation control measures, including silt fences and other barriers. The ESC Plan should be reviewed and updated, if needed, by the chosen construction contractor. The final plan should be developed based on the guidance provided in the *Erosion and Sediment Control Guideline for Urban Construction* (GGHCA 2006). Basic elements of the plan should include consideration of:

- Requirements and timing for rehabilitation of disturbed areas;
- Stormwater management strategies during construction;
- Erosion prevention measures (e.g., hydroseeding, sodding, erosion control matting, tarping of stockpiles);
- Sedimentation control measures (e.g., silt fences and other barriers); and
- Inspection and performance monitoring requirements and adaptive management considerations.

Implementation of an effective ESC Plan, incorporating both erosion and sedimentation controls, coupled with regular inspection and performance monitoring and implementation of any remedial actions necessary to ensure effective performance, is anticipated to be largely effective in preventing the movement of eroded soil particles off-site towards the downstream features north of Hunsden Sideroad.

However, should erosion and off-site sedimentation occur during the construction process, the proposed setbacks from drainage feature on site, which is located more than 50 m from the proposed development within the woodland community will assist in mitigating potential effects on fish and their habitat further downstream within receiving features.

Overall, no adverse effects on the downstream features are predicted to occur as a result of erosion and sedimentation during construction, provided an effective ESC Plan, including monitoring and adaptive management, is implemented.

#### Stormwater Management During Construction

Increases in stormwater runoff from the disturbed areas of the construction site or pumping of groundwater from excavations, potentially resulting in higher flows to the downstream features could cause increases in bed and bank erosion, aside from the obvious potential increase in erosion from the work area.

It is recommended that the contractor consider management of stormwater throughout the construction period as part of the overall ESC Plan, since stormwater flows through disturbed areas are one of the primary causes of erosion and sedimentation from construction sites. Increased volumes of runoff during construction could also potentially result in increases in erosion due to overland flow, particularly if stormwater runoff from the construction area is concentrated. To mitigate these potential effects, stormwater management techniques should

be implemented prior to construction to control surface water runoff throughout the construction period. Implementation of an effective stormwater control plan during construction is anticipated to prevent adverse effects on the watercourse and fish habitat.

Pumping of groundwater from excavations may be required, depending on the depth of the excavation and groundwater level at the time. If pumping is necessary, consideration should be given to the discharge location, and potential impacts on surface water quality and quantity. Mitigation (e.g., sedimentation filter bags) should be provided to ensure that discharge quality criteria are met (e.g., highly turbid water is not discharged to the environment), and mitigation (e.g., rip rap pad) employed to ensure that discharge water does not erode the soils at the immediate discharge location. Implementation of effective mitigation is anticipated to prevent adverse effects on the downstream features and associated fish habitats.

#### Accidental Spills

Accidental spills of potentially hazardous materials (e.g., fuel and oil from heavy equipment), if transported to the watercourse on the Subject Lands, could cause stress or injury to fish, amphibians and other aquatic biota (e.g., benthic invertebrates).

In order to mitigate the potential for adverse effects on these species and their habitats due to accidental spills during construction, it is recommended that the contractor prepare a spill prevention and response plan to outline the material handling and storage protocols, mitigation measures (e.g., spill kits on-site), monitoring measures and spill response plans (i.e., emergency contact procedures, including MOECC Spills Action Centre, and response measures including containment and clean-up). Implementation of an effective spill prevention and response plan is anticipated to be largely effective in preventing adverse effects on these species and their habitats.

## 7.2.2 Potential Post-Construction Impacts

No direct impacts on the downstream features and associated fish habitat are anticipated to occur during the post-construction period, since there would be no requirement for any activity within these features.

However, potential indirect impacts may occur, including:

- Changes in flow and water quality due to stormwater management and changes in groundwater infiltration; and
- Effects on water quality associated with runoff from urban areas.

These potential impacts and recommended mitigation measures is discussed in the following sections.

#### 7.2.2.1 Stormwater Management and Changes in Infiltration

The proposed development and associated changes in soil permeability, and storm water treatment and flows, may affect the flows to the receiving features downstream.

The stormwater management system has been designed to convey water to the same areas as at present, and to relevant regulatory requirements. Compliance with these limits is anticipated to be largely effective at mitigating potential impacts on the downstream features. Consideration should also be given to ensure water balance into the groundwater table is maintained.

Implementation of these measures would mitigate potential impacts on the downstream features and associated fish habitat.

#### 7.2.2.2 Impacts on Water Quality

The proposed stormwater management system has been designed to provide water quality control.

Some surface water on the Subject Lands will infiltrate through residential lawns and into the shallow groundwater or will flow directly as overland runoff from residential rear yards towards the drainage features. This runoff or infiltration water could potentially be impaired due to residential use of potential contaminants (e.g., lawn fertilizers) or other residential land use activities (including accidental spills in rear yards). As the receiving features are situated well away from the proposed development, potential effects on water quality within the receiving features are not anticipated.

# 7.3 Significant Woodlands/Habitat of Endangered and Threatened Species/Confirmed and Candidate Significant Wildlife Habitat

The woodland community and associated confirmed and candidate significant wildlife habitats are being protected from direct impacts (i.e., avoidance) and from indirect impacts, in part through the implementation of the 30 m development setback.

Noise from construction activities may result in wildlife avoidance of the edges abutting active work areas during the construction period. Where possible, construction activities should be timed outside of the nighttime and early morning periods during peak of the bat maternity roosting and bird breeding seasons (typically May through July). Some localized movement of wildlife out of these edge areas may still occur during the construction phase. Given continued development in this area and active agricultural operations, wildlife likely have some tolerance to background noise and so would be somewhat tolerant of construction activities.

Following construction, increased noise in vicinity of the woodland community due to residential activities (e.g., lawn mowing, vehicle movement, etc.), and the potential for increased predation pressure from domestic cats allowed to roam free outdoors may occur. These effects are already present to a degree given the existing residential development to the southwest, as well as rural residential dwellings on the Subject Lands and surrounding properties. Measurable alterations in wildlife composition within the significant woodlands are not anticipated following occupation of the residential subdivision. These potential effects may be further reduced through the development and distribution of a homeowner's manual that explains the relationship between the development and adjacent significant natural areas.

#### 7.4 PNAC Woodland

Though generally of low ecological function, one indirect impact to this PNAC Woodland will be the change in site drainage as a result of the proposed development. Currently, some exiting overland flow is directed towards the pine plantation. Post-development, overland flow will be directed toward grass swales which will drain water into the proposed bioswale along Street A (C.F. Crozier & Associates Inc., 2022) and away from the woodland.

The cultural plantation is a mature community with generally limited ground cover outside of grass species. Pine plantations are tolerant of a range of moisture regimes, and as a result, this minor alteration in hydrology is not expected to impact the feature.

## 7.5 Migratory Birds/Endangered Species of Bats

To ensure that migratory birds or endangered species of bats are not impacted during construction, any vegetation removal or significant earthworks should occur outside of the breeding bird window of April 1 – August 31 (dates approximate) and tree removals should occur outside of the active bat window (April 1 through September 30). In rare circumstances where this window cannot be avoided, a nest search or bat exit survey (as appropriate) is recommended to ensure that the vegetation being removed is not providing active nesting/roosting habitat for these species. If activity is confirmed, a buffer will be marked off surrounding any active nests/roost trees that must be maintained until activity in the nest/roost tree has ceased.

#### 7.6 Air Pollution

Consideration was given to whether there would be an increase in air pollution as a result of the proposed development. There may be a minor increase in air pollution during the construction phase associated with earth moving activities or use of other gas-powered equipment, given the relatively small scale of the development, and nature of the proposed works, these impacts would be anticipated to be short-term and temporary. No measurable impact on local air pollution is anticipated following construction given the low density nature of the proposed development.

## 8. Environmental Management/Reforestation Plan

As identified within section 7 above, and as shown on Figure 4, the following environmental protection measures have been proposed for the natural heritage features surrounding the proposed development:

- 30 m development setback from the significant woodlands/significant wildlife habitats.
   This will consist of a 20 m vegetated buffer and a 10 m portion of the developments lots outside of the structural envelope;
- 20 m vegetated buffer along the existing hedgerow linkage connection between the two significant woodlands; and
- 10 m development setback from the PNAC woodland west of the Subject Lands that will be situated within the developments lots outside of the structural envelope.

Areas situated within the development lots (Lots 3, 4 and 6 to 12) but outside of the structural envelope adjacent to natural heritage features, as well as Block 18, will be seeded with a native upland seed mix. Cover crops should be incorporated to support establishment of the planted seed.

A reforestation planting plan will be prepared at detailed design for the 1.5 ha Block 16 that includes the 20 m vegetated buffer for the significant woodlands and linkage connections, as well as the 0.8 ha Block 17 situated west of the southern significant woodland to the western property line. The reforestation planting plan will be prepared to target extensions of the existing native Fresh-Moist Sugar Maple Hardwood Deciduous Forest (FOD6-5) within the retained significant woodland communities.

As part of the Reforestation Planting Plan prepared at the detail design, the plan will also assess and determine what invasive species management for the Category 1 species is recommended on the Subject Lands. The Category 1 plants observed on the Subject Lands are:

- European Swallowwort;
- Garlic Mustard;
- Dame's Rocket:
- Tartarian Honeysuckle;
- European Buckthorn; and
- Manitoba Maple

A species-specific management assessment will be undertaken to determine whether active management is warranted (i.e., using chemical, biological, or physical interventions) or whether indirect management is appropriate (i.e., supporting natural succession). Consideration will also be given to the likelihood of success in eliminating the target species given occurrence within the local landscape.

#### **Species Selection**

Species selection will consider specific moisture, soil and sun requirements of the area. Some species (e.g., Ash, Elm) will not be selected due to pest and disease concerns that could impact their survivability. Native plant materials should be sourced from native plant nurseries and seed suppliers within 100 km of the Subject Lands, if possible, to reduce transplant shock. All plant materials will be obtained and installed in accordance with the Canadian Nursery Stock Standard. Native shrub and tree species will be selected to provide a diverse assemblage of plant species. Plantings will include fast-growing and pioneer species more tolerant of harsher/variable growing conditions.

The type of planting stock is dependent on the species and their modes of reproduction, as well as practicality. The following plant stock will be considered within the planting areas:

- Herbs (forbs, graminoids): seeds, plugs;
- Shrubs: 1-gallon pots, stem cuttings, rootstock cuttings; and
- Trees: seed, bareroot, ball and burlap, whips, potted seedlings.

#### Soil Amendments

Currently, the lands subject to the reforestation plan are under active agricultural use. Soil testing, prior to planting, to determine the need for soil amendments for healthy plant growth is recommended. Ahead of planting, site preparation is key to ensure that soil moisture capacity and nutrient content are suitable for native plant growth. Native plants generally require low soil nitrogen content and nutrient supplementation is not expected (generally nutrient levels are high in recently farmed cash crop fields). The addition of targeted mycorrhizal inoculants to establish symbiotic relationship between a specific plant and mycorrhiza symbiont (i.e., arbuscular mycorrhiza and Maple (*Acer* ssp.) trees; ectomycorrhizal and Oak (*Quercus* spp.) trees) is generally helpful to facilitate native plant establishment (Chen et al 2016). Studies have also shown that mycorrhizal symbionts are expected to improve the host tree's adaptation to stressors associated with climate change such as high temperatures, drought, salinity and flooding (Usman et al. 2021).

Excessive application of soil amendments (i.e., fertilizers) could negatively impact the surrounding landscape as it will result in nutrient loading and could impact the realigned watercourses. In accordance with CH's Guidelines for Landscaping and Rehabilitation Plans (2021), should soil amendments be required, soil amendments will be sourced from sustainable practices (e.g., incorporating leaf mulch or compost that meets Category AA or A of the MECP Ontario Compost Standard Quality). The incorporation of peat moss is strongly discouraged and composted leaves used instead. Upland disturbed areas should have at least 20 cm of topsoil containing 5 to 15% organic matter (by dry weight) depending on the type of vegetation to be established, a total uncompacted soil depth of at least 30 cm and a soil pH of 6.0 to 8.0 per the TRCA's (2012) Preserving and Restoring Healthy Soil: Best Practices for Urban Construction. Topsoil requirements will also follow CH's (2021) Guidelines for Landscaping and Rehabilitation Plans. Where soil has been compacted, a minimum of 45 cm of clean topsoil will be evenly distributed throughout the site. Imported soil will be mixed with native soil to ensure soil micro-organisms are adapted to the site.

Within tree pits (areas where trees will be planted), trees should have a topsoil layer of a minimum depth of 60 cm. The subsoil layer should be either tilled, scarified or excavated and replaced to a minimum depth of 30 cm. Incorporation of the upper layer into the sublayer should be included to avoid stratified layers where possible. This will produce a total of 90 cm of uncompacted soil, per the TRCA (2012).

All landscaped works, within the reforestation areas, will be reviewed weekly during the construction period to ensure all planting and surface treatments are installed per specifications. It is also anticipated that the works will be inspected with the Town once substantially complete. An additional inspection will be arranged once a year for the two-year compliance period following implementation to ensure that all works are established.

#### 9. Conclusions and Recommendations

This Scoped EIS has been developed as part of the planning process for the proposed development at 10249 Hunsden Sideroad in the Town of Caledon, Ontario. An assessment of impacts on natural features and their associated functions has been conducted and discussed in relation to the PPS and the ORMCP.

The proposed development occurs in areas that are agricultural, and a 30 m development setback has been proposed from the natural heritage features associated with the significant woodlands, while a 20 m setback for the linkage corridor, and a 10 m setback for the PNAC woodland is provided.

Based upon the natural heritage feature inventories and analyses carried out, the following conclusions are drawn:

- The agricultural lands upon which development is proposed, do not provide habitat for any significant natural features/KNHF/KHF when considered through both the PPS and the ORMCP;
- The significant woodlands on the Subject Lands were also identified as providing habitat for endangered species (i.e., Butternut, bat SAR), and have also been identified as candidate and confirmed significant wildlife habitat:
  - o Candidate Bat Maternity Colonies (both FODs);
  - o Candidate Reptile Hibernacula (south-most FOD);
  - Confirmed Woodland Area-Sensitive Bird Breeding Habitat (south-most FOD);
  - Confirmed Habitat for Species of Conservation Concern (both FODs):
    - Eastern Wood-Pewee; and
    - Wood Thrush.
- A PNAC woodland has been identified to the west of the Subject Lands that is of low ecological function;
- The identified development setbacks will sufficiently protect the identified natural heritage features;
- Linkage functions along the eastern extent of the Subject Lands will be maintained. Should a roadway be required to cross to the lands to the East, careful consideration should be given to ensuring appropriate design as this area is likely to see greater wildlife traffic given the proximity between two significant features on the landscape; and
- A formal Erosion and Sediment Control Plan will be provided as part of the detailed design phase of the Project. The use of standard mitigation measures regarding the use of fuels and chemicals during the construction process will reduce the risk of groundwater or surface water contamination from accidental spills.

Based upon current and available technical information and analyses, the predicted effects on the natural features and associated functions will be avoided/minimized through the protection, mitigation and enhancement measures recommended and discussed in this report.

#### Prepared By:

**GEI Consultants** 

Julie Snow Ecologist 249.877.8486

jsnow@geiconsultants.com

Sean Male Project Manager 289.407.7483

smale@geiconsultants.com

## **References and Background Materials**

Bird Studies Canada, Environment Canada's Canadian Wildlife Service, Ontario Nature, Ontario Field Ornithologists and Ontario Ministry of Natural Resources. 2006. Ontario Breeding Bird Atlas Website. http://www.birdsontario.org/atlas/index.jsp

Brouillet, L., F. Coursol, S.J. Meades, M. Favreau, M. Anions, P. Bélisle & P. Desmet 2010+. VASCAN, the Database of Vascular Plants of Canada. http://data.canadensys.net/vascan/

C.F. Crozier & Associates Inc. 2023. Functional Servicing and Preliminary Stormwater Management Report. 10249 Hunsden Sideroad Estate Residential Development. Town of Caledon

Cadman, M.D., D.A. Sutherland, G.G. Beck, D. Lepage, and A.R. Courturier (eds.) 2007. Atlas of the breeding birds of Ontario, 2001-2005. Bird Studies Canada, Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature, Toronto, xxii + 706 pp.

Cadman, M.D., H.J. Dewar, and D.A. Welsh. 1998. The Ontario Forest Bird Monitoring Program (1987-1997): Goals, methods and species trends observed. Technical Report Series No. 325, Canadian Wildlife Service

Chen, W., Koide R.T., Adams T.S., DeForest J.L., Cheng L., and Eissenstat D.M. 2016. Root morphology and mycorrhizal symbioses together shape nutrient foraging strategies of temperate trees. National Academy of Sciences. 113(31) 8741 – 8746.

Conservation Halton (CH). 2021. Guidelines for Landscaping and Rehabilitation Plans. Available online at: https://www.conservationhalton.ca/policies-and-guidelines

Department of Fisheries and Oceans (DFO). 2023. Aquatic Species at Risk Distribution Mapping. Available online at http://www.dfo-mpo.gc.ca/species-especes/sara-lep/map-carte/index-eng.html

eBird. 2023. eBird: An online database of bird distribution and abundance. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: http://www.ebird.org.

GEI Consultants. 2023. Fluvial Geomorphic Site Visit – Drainage Feature, Tributary of Beeton Creek, 10249 Hunsden Sideroad (GEI PN2101948)

Government of Canada. 1985. Fisheries Act (R.S.C., 1985, c. F-14). (Last Amended August 2019).

Government of Canada. 1994. Migratory Birds Convention Act (S.C. 1994, c. 22). (Last Amended December 2017)



Government of Ontario. 1990. Ontario Regulation 172/06: Nottawasaga Valley Conservation Authority: Regulation of Development, Interference with Wetlands and Alterations to Shorelines and Watercourses under Conservation Authorities Act, 1990, R.S.O. 1990, c.C.27.

Government of Ontario. 2001. Ontario Regulation 140/02: Oak Ridges Moraine Conservation Plan under Oak Ridges Moraine Conservation Act, 2001, S.O. 2001, c.31.

Government of Ontario. 2007a. Endangered Species Act, 2007, S.O. 2007, c. 6. (Consolidated October 2021).

Government of Ontario. 2007b. Ontario Regulation 230/08: Species at Risk in Ontario List. Endangered Species Act, 2007, S.O. 2007, c. 6. (Consolidated August 2018).

Greater Golden Horseshoe Area Conservations Authorities (GGHCA) 2006. Erosion and Sediment Control Guideline for Urban Construction.

iNaturalist. 2023. Available from https://www.inaturalist.org.

Lee, H.T., W.D. Bakowsky, J. Riley, J. Bowles, M. Puddister, P. Uhlig and S. McMurray. 1998. Ecological land classification for Southern Ontario: first approximation and its application. Ontario Ministry of Natural Resources, South Central Region, Science Development and Transfer Branch. Technical Manual ELC-005.

North-South Environmental Inc (NSE), Dougan & Associates and Sorensen Gravely Lowes 2009. Peel-Caledon Significant Woodlands and Significant Wildlife Habitat Study. Report prepared for the Region o Peel and the Town of Caledon, Ontario. xi+187pp+app.

Nottawasaga Valley Conservation Authority (NVCA) 2023. Innisfil Creek Subwatershed Health Check 2023. Available online: https://www.nvca.on.ca/wp-content/uploads/2023/07/Watershed%20Health%20Check\_Innisfil%20Creek%20Subwatershed-opt.pdf

Oldham, M.J., W.D. Bakowsky and D.A. Sutherland. 1995. Floristic quality assessment for southern Ontario. OMNR, Natural Heritage Information Centre, Peterborough. 68 pp.

Ontario Ministry of Municipal Affairs and Housing (MMAH). 2020. Provincial Policy Statement. Ontario Ministry of Municipal Affairs and Housing. Toronto: Queens Printer for Ontario. 37 pp.

Ontario Ministry of Natural Resources (MNR) 2010. Natural Heritage Reference Manual for the Natural Heritage Policies of the Provincial Policy Statement. Available online: http://www.mnr.gov.on.ca/en/Business/LUEPS/Publication/249081.html

Ontario Ministry of Natural Resources (MNR). 2000. Significant Wildlife Habitat Technical Guide. Fish and Wildlife Branch, Wildlife Section, Science Development and Transfer Branch, Southcentral Sciences Section. 151 pp.

Ontario Ministry of Natural Resources and Forestry (MNRF) 2015. Significant Wildlife Habitat Ecoregion 6E Criterion Schedule. Available online at https://dr6j45jk9xcmk.cloudfront.net/documents/4775/schedule-6e-jan-2015-access-ver-final-s.pdf

Ontario Ministry of Natural Resources and Forestry (MNRF). 2024. Natural Heritage Information Centre database. Available online at https://www.ontario.ca/page/get-natural-heritage-information

Ontario Nature. 2020. Ontario Reptile and Amphibian Atlas. Available online at https://ontarionature.org/programs/citizen-science/reptile-amphibian-atlas

Region of Peel Official Plan. 2021, Office Consolidation 2021. Available online at: https://www.peelregion.ca/officialplan/download/

Toronto Entomologists' Association. 2023. Ontario Butterfly Atlas Online. Available online at http://www.ontarioinsects.org/atlas/index.html

Toronto Entomologists' Association. 2020. Ontario Moth Atlas Online. Available online at http://www.ontarioinsects.org/moth/

Toronto and Region Conservation Authority (TRCA). 2012. Preserving and Restoring Healthy Soil: Best Practices for Urban Construction.

Toronto and Region Conservation Authority (TRCA) and Credit Valley Conservation (CVC). 2014. Evaluation, Classification and Management of Headwater Drainage Features Guidelines. Available online at: https://cvc.ca/wp-content/uploads//2021/06/HDFA-final.pdf

Town of Caledon Official Plan. 2018, Office Consolidation 2018. Available online at: https://www.caledon.ca/en/town-services/official-plan.aspx

Usman, M., Ho-Plágaro, T., Frank H.E.R., Calvo-Polanco M., Gaillard I., Garcia K., Zimmermann S.D. 2021. Mycorrhizal Symbiosis for Better Adaptation of Trees to Abiotic Stress Caused by Climate Change in Temperate and Boreal Forests. Frontiers in Forests and Global Change 4 (2021). https://www.frontiersin.org/articles/10.3389/ffgc.2021.742392/full

Varga, S., editor. 2005. Distribution and status of the vascular plants of the Greater Toronto Area. Ontario Ministry of Natural Resources, Aurora District. 96 pp.

Urban Forest Associates Inc. 2002. Invasive Exotic Species Ranking for Southern Ontario. 7pp.

# Appendix A


#### **Figures**

Figure 1 – Project Location and Landscape Context

Figure 2 – Ecological Land Classification

Figure 3 – Designated Natural Heritage Features

Figure 4 – Draft Plan





NOTES:

1. Coordinate System: NAD 1983 UTM Zone 17N.
2. Base features produced under license with the Ontario Ministry of Natural Resources and Forestry © Queen's Printer for Ontario, 2023.
3. Orthoimagery © First Base Solutions, 2023. Imagery taken in 2021.
4. Dripline as staked with the Town of Caledon on August 9, 2023

Subject Lands

Watercourse

Butternut

Butternut 25m Protection Zone

**Ecological Land Classification** 

-- - Dripline4

#### **ELC Legend**

AG, Agricultural

CUP, Cultural Plantation

FOD6-5, Fresh – Moist Sugar Maple Hardwood Deciduous Forest

HR, Hedgerow

RES, Residential


Carringwood Homes 10249 Hunsden Sideroad, Caledon, ON

## Figure 2 **Ecological Land Classification**









Coordinate System: NAD 1983 UTM Zone 17N.
 Base features produced under license with the Ontario Ministry of Natural Resources and Forestry © Queen's Printer for Ontario, 2024.
 Orthoimagery © First Base Solutions, 2024.
 Imagery taken in 2021.

Oak Ridges Moraine Conservation Plan – Natural Linkage Area/Town of Caledon Policy Area 4

Ecological Land Classification Potential Natural areas and Corridors Woodland

Candidate Reptile Hibernacula

Intermittent Watercourse/Fish Habitat

Butternut 25m Protection Zone

---- Candidate Significant Wildlife Habitat (Reptile Hibernacula)

FOD6-5, Fresh – Moist Sugar Maple Hardwood Deciduous Forest


HR, Hedgerow

Figure 3 Designated Natural Heritage Features

1:4,000







Confirmed Significant Wildlife Habitat (Woodland Area-Sensitive Bird Breeding Habita Ecological Land Classification

Candidate Reptile Hibernacula

Butternut

PNAC Woodlands + 10m

Significant Woodland + 30 m


Candidate Significant Wildlife Habitat (Reptile Hibernacula)

Intermittent Watercourse/Fish Habitat

■ ■ ■ Drainage Swale







# **Appendix B**

#### **Tables**

Table 1 - Field Studies and Natural Inventories

Table 2 – Master Plant List

Table 3 - Master Bird

Table 4 – Master Wildlife List

Table 5 – Significant Wildlife Habitat Assessment

Table 6 – Predicted Effects, Mitigation, Enhancement and Net Effects



Table 1: Field Studies and Natural Inventories (2021 - 2022)

SURVEYORS	SURVEY	SURVEY TYPE	DATE	TIM	ИЕ	AIR	WATER	HUMIDITY	CLOUD	BEAUFORT	PRECIPITATION
(SURNAME, INTL)	ROUND		(2022)	START	END	TEMP (C°)	TEMP (C°)	(%)	COVER (%)	WIND SPEED	COMMENTS
					20	21					
Szabo, A	1	Site Reconnaissance	02-JN	-	-	24	N/A	37	60	1	None
					20	22					
Kimble, B.	1	Headwater Drainage Feature Assessment	20-AP	11:00	13:30	4	N/A	54	20	4	None
Snow J.	1	Spring Botanical Inventory and ELC	17-MA	09:00	12:00	14	N/A	62	80	4	None
Kimble, B.	2	Headwater Drainage Feature Assessment	20-MA	11:00	11:30	23	N/A	N/A	70	3	None
Burke, P.	1	Breeding Bird Surveys	14-JN	05:30	07:40	17	N/A	63	5	1	None
Burke, P.	2	Breeding Bird Surveys	30-JN	05:40	07:30	17	N/A	66	85	1	None
Snow J.	2	Summer Botanical Inventory and ELC	12-JL	09:45	12:00	22	N/A	51	45	2-3	None
Snow J.	3	Fall Botanical Inventory and ELC	02-SE	10:00	12:00	23	N/A	54	60	1	None

LEGEND:



#### Table 1: Field Studies and Natural Inventories (2021 - 2022)

	BEAUFORT WIND SPEED SCALE	MON	ITH (CODE)
0	Calm (<1 km/hr)	JA	January
1	Light Air (1-5 km/hr)	FB	February
2	Light Breeze (6-11 km/hr)	MR	March
3	Gentle Breeze (12-19	AP	April
	km/hr)	MA	May
4	Moderate Breeze (20-28	JN	June
	km/hr)	JL	July
	·	AU	August
		SE	September
		OC	October
		NO	November
		DE	December



ORDER	FAMILY	LATIN NAME	COMMON NAME	COEFFICIENT OF CONSERVATISM	WETNESS INDEX	OWES WETLAND SPECIES	WEEDINESS INDEX	INVASIVE EXOTIC RANK (Urban Forest Associates 2002)	PROVINCIAL STATUS (S-RANK)	GLOBAL STATUS (G-RANK)	COSSARO (MNRF)	COSEWIC STATUS	PEEL (Varga 2005)	FOD6-5 (A)	hedgerows
DICOTYLEDONS	Adoxaceae	Sambucus racemosa ssp. pubens	Red Elderberry	5	3			Р	S5	G5			Х	хх	
DICOTYLEDONS	Adoxaceae	Viburnum acerifolium	Maple-Leaved Viburnum	6	5				S5	G5			Х	х	
DICOTYLEDONS	Adoxaceae	Viburnum rafinesquianum	Downy Arrowwood	7	5				S5	G5			R4	x x	. 📙
DICOTYLEDONS	Amaranthaceae	Chenopodium album	Common Lamb's-Quarters		3		-1		SNA	G5T5			Х		х
DICOTYLEDONS	Anacardiaceae	Rhus typhina	Staghorn Sumac	1	3				S5	G5			X	х	х
DICOTYLEDONS	Apiaceae		Poison Hemlock		-3		-1		SNA	G5			X	x x	+
DICOTYLEDONS DICOTYLEDONS	Apiaceae	Daucus carota	Wild Carrot	2	5		-2		SNA	GNR G5T?			X X	x x	x
DICOTYLEDONS	Apocynaceae Apocynaceae	Apocynum androsaemifolium Asclepias syriaca	Spreading Dogbane Common Milkweed	0	5				S5 S5	G5			X	X X	₩
DICOTYLEDONS	Apocynaceae	Vinca minor	Lesser Periwinkle	U	5		-2	2	SNA	GNR			X	^ x	+
DICOTYLEDONS	Apocynaceae		European Swallowwort		5		-	1	SNA	GNR			X	x x	
DICOTYLEDONS	Aristolochiaceae	Asarum canadense	Canada Wild-Ginger	6	5			_	S5	G5			X	x x	
DICOTYLEDONS	Asteraceae	Ambrosia artemisiifolia	Common Ragweed	0	3				S5	G5			Х	х	х
DICOTYLEDONS	Asteraceae		Common Burdock		3		-2		SNA	G?T?			Х	x x	. х
DICOTYLEDONS	Asteraceae	Cirsium vulgare	Bull Thistle		3		-1		SNA	G5			Х		х
DICOTYLEDONS	Asteraceae	Erigeron annuus	Annual Fleabane	0	3				S5	G5			Х	x x	x
DICOTYLEDONS	Asteraceae	Euthamia graminifolia	Grass-Leaved Goldenrod	2	0				S5	G5			Х		х
DICOTYLEDONS	Asteraceae		Black-Eyed Susan	0	3				S5	G5			Х	х	
DICOTYLEDONS	Asteraceae		Tall Goldenrod	1	3				S5	GNR			X	x	x
DICOTYLEDONS	Asteraceae	Solidago canadensis	Canada Goldenrod	1	3				S5	G5T5			X	+	х
DICOTYLEDONS	Asteraceae	Solidago gigantea	Giant Goldenrod	4	-3	Т			S5	G5		<del>                                     </del>	X	x x	₽
DICOTYLEDONS	Asteraceae		Grey-Stemmed Goldenrod (var. nemor	2	5				SS SNA	G5T?			X X	++	X
DICOTYLEDONS	Asteraceae	Sonchus arvensis ssp. arvensis	Field Sow-Thistle	5	3				SNA	GNR				х	х
DICOTYLEDONS DICOTYLEDONS	Asteraceae Asteraceae	Symphyotrichum cordifolium Symphyotrichum ericoides var. ericoide	Heart-Leaved Aster	5	5				S5 S5	G5 G5T5		<del>                                     </del>	X	×	₩
DICOTYLEDONS	Asteraceae		Panicled Aster	3	-3	1			SS SS	G515 G5		<del>                                     </del>	X	x	x
DICOTYLEDONS	Asteraceae	Symphyotrichum novae-analiae		2	-3				S5	G5		-	X	x	x
DICOTYLEDONS	Asteraceae	Taraxacum officinale	New England Aster Common Dandelion		-3		-2		SNA	G5	-		X	хх	
DICOTYLEDONS	Asteraceae	Tragopogon dubius	Yellow Goatsbeard		5		-1		SNA	GNR			X	^ ^	+
DICOTYLEDONS	Asteraceae		Coltsfoot		3	т	-2	4	SNA	GNR			X	· ·	-
DICOTYLEDONS	Balsaminaceae		Spotted Jewelweed	4	-3	i	-	,	S5	G5			X	x x	+
DICOTYLEDONS	Berberidaceae	Caulophyllum thalictroides	Blue Cohosh	5	5				S5	G5			R1	x x	
DICOTYLEDONS	Berberidaceae	Podophyllum peltatum	May-Apple	5	3				S5	G5			X	х х	+
DICOTYLEDONS	Betulaceae	Betula alleghaniensis	Yellow Birch	6	0	T			S5	G5			Х	хх	$\top$
DICOTYLEDONS	Betulaceae	Betula papyrifera	Paper Birch	2	3	T			S5	G5			Х	х	$\top$
DICOTYLEDONS	Betulaceae	Ostrya virginiana	Eastern Hop-Hornbeam	4	3				S5	G5			Х	хх	$\top$
DICOTYLEDONS	Boraginaceae	Hackelia virginiana	Virginia Stickseed	5	3				S5	G5			U	x x	.
DICOTYLEDONS	Boraginaceae	Hydrophyllum virginianum var. virginia	Virginia Waterleaf	6	0				S5	G5			Х	x x	. х
DICOTYLEDONS	Boraginaceae	Symphytum officinale	Common Comfrey		5		-1		SNA	GNR			Х	хх	. 🔲
DICOTYLEDONS	Brassicaceae	Alliaria petiolata	Garlic Mustard		0		-3	1	SNA	G5			Х	x x	!
DICOTYLEDONS	Brassicaceae	Cardamine diphylla	Two-Leaved Toothwort	7	3				S5	G5			Х	х	
DICOTYLEDONS	Brassicaceae	Hesperis matronalis	Dame's Rocket		3		-3	1	SNA	G4G5			Х	х	х
DICOTYLEDONS	Brassicaceae		Field Pennycress		5		-1		SNA	GNR			Х		х
DICOTYLEDONS	Campanulaceae	Lobelia inflata	Indian-Tobacco	3	3				S5	G5			Х	х	
DICOTYLEDONS	Caprifoliaceae	Lonicera canadensis	Canada Fly Honeysuckle	6	3		_		S5	G5			X	x	+
DICOTYLEDONS	Caprifoliaceae	Lonicera tatarica	Tartarian Honeysuckle		3		-3 -3	1	SNA SNA	GNR GNR			X X	х	x
DICOTYLEDONS DICOTYLEDONS	Caryophyllaceae		Bouncing-Bet		3 5		-3 -1	3	SNA	GNR			X		
	Caryophyllaceae Cornaceae	Silene vulgaris Cornus alternifolia	Bladder Campion	6	3		-1		SS SS	GNR G5		-	X	x x	X
DICOTYLEDONS DICOTYLEDONS	Cucurbitaceae	Echinocystis lobata	Alternate-Leaved Dogwood Wild Cucumber	3	-3	Т		1	55 S5	G5		<del>                                     </del>	X	-	+
DICOTYLEDONS	Elaeagnaceae		Russian Olive	3	-3	-	-1	3	SNA	GNR			X	++	x
DICOTYLEDONS	Fabaceae		American Hog-Peanut	4	0	Т	-	,	S5	G5			X	×	+
DICOTYLEDONS	Fabaceae	Lotus corniculatus	Garden Bird's-Foot Trefoil		3		-2	2	SNA	GNR			X	T	х
DICOTYLEDONS	Fabaceae	Medicago sativa ssp. sativa	Alfalfa (ssp. sativa)		5		-1	4	SNA	GNRTNR			X		x
DICOTYLEDONS	Fabaceae		White Sweet-Clover		3		-3	2	SNA	GNR			Х		х
DICOTYLEDONS	Fabaceae	Trifolium pratense	Red Clover		3		-2	4	SNA	GNR			Х	х	х
DICOTYLEDONS	Fabaceae	Vicia cracca	Tufted Vetch		5		-1	2	SNA	GNR			Х		х
DICOTYLEDONS	Fagaceae	Fagus grandifolia	American Beech	6	3				S4	G5			Х	хх	╜
DICOTYLEDONS	Geraniaceae	Geranium robertianum	Herb-Robert	2	3		-2		S5	G5			Х	x x	╜
DICOTYLEDONS	Grossulariaceae	Ribes cynosbati	Eastern Prickly Gooseberry	4	3				S5	G5			Х	x x	┷-
DICOTYLEDONS	Grossulariaceae	Ribes rubrum	European Red Currant		5	T	-2		SNA	G4G5			X	x x	╨
DICOTYLEDONS	Juglandaceae	Juglans cinerea	Butternut	6	3				S2?	G4	END	END	Х	х	$\perp \!\!\! \perp \!\!\! \perp$
DICOTYLEDONS	Juglandaceae		Black Walnut	5	3				S4?	G5			X	x	х
DICOTYLEDONS	Lamiaceae	Ajuga reptans	Creeping Bugleweed	+	5		-1	4	SNA	GNR			X	х	
DICOTYLEDONS	Lamiaceae	Leonurus cardiaca ssp. cardiaca	Common Motherwort		5		-2		SNA	GNR			X	x x	
DICOTYLEDONS	Malvaceae	Tilia americana	Basswood	4	3				S5	G5	-	-	X	x x	
DICOTYLEDONS DICOTYLEDONS	Oleaceae	Fraxinus americana	White Ash	3	-3	т			S4 S4	G5 G5	-	-	X	x x	
	Oleaceae	,	Red Ash	3	-3 5	I	2	2						X X	ŧIJ
DICOTYLEDONS DICOTYLEDONS	Oleaceae		Common Lilac Canada Enchanter's Nightshade	2	3		-2	2	SNA S5	GNR G5T5			X X	x x	x
DICOTYLEDONS	Onagraceae Oxalidaceae	Circaea canadensis ssp. canadensis Oxalis stricta		0	3				S5 S5	G515 G5			X	x x	
DICOTYLEDONS			European Wood-Sorrel Squirrel-Corn	7	5				55 55	G5	-	<del>                                     </del>	X U	x x	
DICOTYLEDONS	Papaveraceae Papaveraceae	Sanguinaria canadensis	Bloodroot	5	3				S5	G5			X	X X	
DICOTYLEDONS	Phrymaceae	Phryma leptostachya var. leptostachya		6	3				S4S5	G5	-		X	×	+
DICOTYLEDONS	Plantaginaceae	Linaria vulgaris	Butter-And-Eggs	Ü	5		-1	4	SNA	GNR	-		X		y
	Plantaginaceae		English Plantain		3		-1	T	SNA	G5			X	11	x
DICOTYLEDONS				1										-	لت
DICOTYLEDONS DICOTYLEDONS	Plantaginaceae		Common Plantain		3		-1		SNA	G5			X	1 1	x



ORDER	FAMILY	LATIN NAME	COMMON NAME	COEFFICIENT OF CONSERVATISM	WETNESS INDEX	OWES WETLAND SPECIES	WEEDINESS INDEX	INVASIVE EXOTIC RANK (Urban Forest Associates 2002)	PROVINCIAL STATUS (S-RANK)	GLOBAL STATUS (G-RANK)	COSSARO (MNRF)	COSEWIC STATUS	PEEL (Varga 2005)	FOD6-5 (A) FOD6-5 (B)	hedgerows
DICOTYLEDONS	Polygonaceae	Rumex crispus	Curled Dock		0	T	-2		SNA	GNR			Х	x x	
DICOTYLEDONS	Primulaceae	Lysimachia borealis	Northern Starflower	6	0				S5	G5T?			Х	х	
DICOTYLEDONS	Ranunculaceae	Actaea pachypoda	White Baneberry	6	5				S5	G5			Х	x x	
DICOTYLEDONS	Ranunculaceae	Actaea rubra ssp. rubra	Red Baneberry	6	3				S5	G5			Х	x x	
DICOTYLEDONS	Ranunculaceae	Anemone acutiloba	Sharp-Lobed Hepatica	8	5				S5	G5			X	x	44
DICOTYLEDONS DICOTYLEDONS	Ranunculaceae Ranunculaceae	Anemone virginiana Ranunculus abortivus	Tall Anemone Kidnev-Leaved Buttercup	2	0				S5 S5	G5 G5			X X	x x	х
DICOTYLEDONS	Ranunculaceae	Thalictrum dioicum	Early Meadow-Rue	6	3				55 55	G5			X	X X	+
DICOTYLEDONS	Rhamnaceae	Rhamnus cathartica	European Buckthorn	0	0	Т	-3	1	SNA	GNR			X	x x	+
DICOTYLEDONS	Rosaceae	Fragaria virginiana	Wild Strawberry	2	3		- 5	-	S5	G5			X	- ^ ^	+^-
DICOTYLEDONS	Rosaceae	Geum aleppicum	Yellow Avens	2	0	T			S5	G5			X	x	Ħ
DICOTYLEDONS	Rosaceae	Geum urbanum	Wood Avens		5		-1		SNA	G5			Х	x	Ħ
DICOTYLEDONS	Rosaceae	Prunus nigra	Canada Plum	4	3				S4	G4G5			U	х х	T
DICOTYLEDONS	Rosaceae	Prunus pensylvanica	Pin Cherry	3	3				S5	G5			Х	х х	
DICOTYLEDONS	Rosaceae	Prunus virginiana var. virginiana	Chokecherry	2	3				S5	G5T?			Х	х	х
DICOTYLEDONS	Rosaceae	Rubus allegheniensis	Alleghany Blackberry	2	3				SU	G5			Х	x	
DICOTYLEDONS	Rosaceae	Rubus idaeus ssp. idaeus	European Red Raspberry		3				SNA	G5T5				x x	х
DICOTYLEDONS	Rosaceae	Rubus occidentalis	Black Raspberry	2	5				S5	G5			X	x x	
DICOTYLEDONS	Rubiaceae	Galium aparine	Common Bedstraw	4	3			-	S5	G5			R4	x x	+
DICOTYLEDONS	Rubiaceae	Galium triflorum	Three-Flowered Bedstraw	4	3	-		-	S5	G5	-		X	x	늰
DICOTYLEDONS	Salicaceae	Populus tremuloides	Trembling Aspen	0	0	T		-	S5	G5			X X	x	X
DICOTYLEDONS	Sapindaceae	Acer negundo	Manitoba Maple		0	T		1	S5	G5					х
DICOTYLEDONS	Sapindaceae	Acer saccharum	Sugar Maple	6	-5	1		-	SS SNA	G5	-		X	x x	
	Sapindaceae	Acer x freemanii	Freeman's Maple	•	-5	T				GNA			XSK	×	х
DICOTYLEDONS DICOTYLEDONS	Saxifragaceae Scrophulariaceae	Tiarella cordifolia  Verbascum thapsus ssp. thapsus	Heart-Leaved Foamflower Common Mullein	6	5	ı	-2		SS SNA	G5 GNR			X		х
DICOTYLEDONS	Solanaceae	Solanum dulcamara	Bittersweet Nightshade		0	Т	-2	3	SNA	GNR			X		x
DICOTYLEDONS	Ulmaceae	Ulmus americana	White Elm	3	-3	T	-2	3	SS SS	G5			X	x x	
DICOTYLEDONS	Urticaceae		European Stinging Nettle	3	0	1	-1	3	SNA	G5T5?			XSR	x x	
DICOTYLEDONS	Violaceae	Urtica dioica ssp. dioica Viola canadensis var. canadensis	Canada Violet	6	3		-1		S5	GNR			X	x x	
DICOTYLEDONS	Violaceae	Viola pubescens	Downy Yellow Violet	5	3				S5	G5			X	x x	-
DICOTYLEDONS	Vitaceae	Parthenocissus quinquefolia	Virginia Creeper	6	3				S4?	G5			RLR	x x	
DICOTYLEDONS	Vitaceae	Parthenocissus vitacea	Thicket Creeper	4	3				S5	G5			X	x x	
DICOTYLEDONS	Vitaceae	Vitis riparia	Riverbank Grape	0	0				S5	G5			Х	хх	х
GYMNOSPERMS	Pinaceae	Picea pungens	Blue Spruce		3				SNA	G5					х
GYMNOSPERMS	Pinaceae	Pinus strobus	Eastern White Pine	4	3	T			S5	G5			Х	х х	
GYMNOSPERMS	Pinaceae	Pinus sylvestris	Scots Pine		3		-3	2	SNA	GNR			Х		х
GYMNOSPERMS	Pinaceae	Tsuga canadensis	Eastern Hemlock	7	3	T			S5	G5			Х	хх	
	Amaryllidaceae	Allium tricoccum var. tricoccum	Wild Leek	7	3				S4	G5			Х	х	
MONOCOTYLEDONS	Araceae	Arisaema triphyllum ssp. triphyllum	Jack-In-The-Pulpit	5	-3	T			S5	G5			Х	x x	
MONOCOTYLEDONS	Asparagaceae	Convallaria majalis var. majalis	European Lily-Of-The-Valley		5		-2	3	SNA	G5			Х	х	$\perp$
MONOCOTYLEDONS	Asparagaceae		Wild Lily-Of-The-Valley (ssp. canadense	5	3				S5	G5T5			Х	x x	$\perp$
MONOCOTYLEDONS	Asparagaceae	Maianthemum racemosum	Large False Solomon's Seal	4	3				S5	G5T			Х	x x	
	Asparagaceae	Polygonatum biflorum var. commutatu		8	3				S4	G5				x x	$\perp$
	Asparagaceae	Polygonatum pubescens	Hairy Solomon's Seal	5	5				S5	G5			X	x	
	Cyperaceae	Carex albursina	White Bear Sedge	7 5	5				S5	G5			R10	x x	
	Cyperaceae	Carex communis yar, communis	Drooping Woodland Sedge	6	5			+	S5	G5?			X X	x x	
MONOCOTYLEDONS MONOCOTYLEDONS	Cyperaceae Cyperaceae	Carex communis var. communis Carex intumescens	Fibrous-Root Sedge Bladder Sedge	6	-3			+	S5 S5	G5 G5			X	A X	+
MONOCOTYLEDONS	Cyperaceae	Carex Intumescens Carex laxiflora	Loose-Flowered Sedge	5	0				S5	G5	-		^ R7	x ^	$\forall$
MONOCOTYLEDONS	Cyperaceae	Carex leptonervia	Finely-Nerved Sedge	5	0	1	I .		S5	G5	1		R6	x x	$\forall$
	Cyperaceae	Carex pensylvanica	Pennsylvania Sedge	5	5				S5	G5			X	x x	
	Cyperaceae	Carex sprengelii	Sprengel's Sedge	6	0				S5	G5			R1	x	
	Cyperaceae	Carex sylvatica	European Woodland Sedge		3		-1		SNA	GNR				x x	П
	Liliaceae	Erythronium americanum ssp. america		5	5				S5	G5T5			Х	x x	
MONOCOTYLEDONS		Trillium erectum	Red Trillium	6	3				S5	G5			Х	x x	
	Melanthiaceae	Trillium grandiflorum	White Trillium	5	3				S5	G5			Х	x x	ш
MONOCOTYLEDONS	Poaceae	Bromus inermis	Smooth Brome		5		-3	4	SNA	G5TNR			Х		х
MONOCOTYLEDONS	Poaceae	Dactylis glomerata	Orchard Grass		3		-1	3	SNA	GNR			Х	х	х
MONOCOTYLEDONS	Poaceae	Phalaris arundinacea var. arundinacea		0	-3	T		Р	S5	GNR	1		Х	$\perp \perp$	х
MONOCOTYLEDONS	Poaceae	Phleum pratense ssp. pratense	Common Timothy		3		-1		SNA	GNR			Х	$\perp$	х
	Smilacaceae	Smilax herbacea	Herbaceous Carrionflower	5	0			-	S4?	G5			X	x x	+
	Xanthorrhoeaceae	Hemerocallis fulva	Orange Daylily		5		-3	4	SNA	GNR			X	x	+
	Xanthorrhoeaceae	Hemerocallis Iilioasphodelus	Yellow Daylily		5	-	-1	-	SNA	GNR			X	+	Х
	Athyriaceae	Athyrium filix-femina var. angustum	Northeastern Lady Fern	4	0	T		-	S5	G5T5	-	-	X	х	+
PTERIDOPHYTES PTERIDOPHYTES	Cystopteridaceae Dennstaedtiaceae	Cystopteris species	Bladder Fern Species Eastern Bracken Fern	5 2	-3 3	Т		-	S5 S5	G5 G5T	-		X	x x	+
PTERIDOPHYTES	Dryopteridaceae	Pteridium aquilinum var. latiusculum Dryopteris intermedia	Evergreen Wood Fern	5	0			<b>-</b>	S5 S5	G51	-	-	X	x x	+
PTERIDOPHYTES	Dryopteridaceae	Polystichum acrostichoides	Christmas Fern	5	2			<b>-</b>	S5 S5	G5	-	-	X	X X	+
PTERIDOPHYTES	Onocleaceae	Matteuccia struthiopteris var. pensylva		5	0	T		1	S5 S5	G5			X	x x	+
PTERIDOPHYTES	Osmundaceae	Osmunda claytoniana	Interrupted Fern	7	0	T			S5	G5			^ R7	x x	
	Pteridaceae	Adiantum pedatum	Northern Maidenhair Fern	7	3				S5	G5	-		X	X X	$\forall$
		riorariani pedatani			, ,		l	1		- 65	1		^	1^1^	لــــــــــــــــــــــــــــــــــــــ

INVASIVE EXOTIC

RANK

WEEDINESS INDEX

PROVINCIAL

STATUS (S-RANK)

GLOBAL STATUS COSSARO

(MNRF)

(G-RANK)

COSEWIC

STATUS

PEEL (Varga 2005)



							_
ORDER	FAMILY	LATIN NAME	COMMON NAME	COEFFICIENT OF CONSERVATISM	WETNESS INDEX	OWES WETLAND SPECIES	
	STATISTICS						
	Species Diversity						
	Total Number of Species:	150					
	Native Species:	101	67%				
	Exotic Species:	49	33%				
	S1-S3 Species:	1	1%				
	S4 Species:	10	10%				
	S5 Species:	89	88%				
	Floristic Quality Assessment (FQA)						
	Mean Co-efficient of Conservatism (CC)	4.2					
	CC 0 - 3 = lowest sensitivity	32	32%				
	CC 4 - 6 = moderate sensitivity	60	59%				
	CC 7 - 8 = high sensitivity	10	10%				
	CC 9 - 10 = highest sensitivity	0	0%				
	Floristic Quality Index (FQI)	42					
	Weedy & Invasive Species						
	Mean Weediness Index (Oldham et al):	-1.7					
	-1 = low potential invasiveness	21	43%				
	-2 = moderate potential invasiveness	15	31%				
	-3 = high potential invasivenss	9	18%				
	Mean Exotic Rank (Urban Forest Associates):	3					
	Category 1	6	12%				
	Category 2	6	12%				
	Category 3	6	12%				
	Category 4	7	14%				
	Potentially Invasive (P)	2	4%				
	Wetland Species						
	Mean Wetness Index	2.5					
	Upland	38	25%				
	Facultative upland	76	51%				
	Facultative	23	15%				
	Facultative wetland	12	8%				
	Obligate wetland	1	1%				
	9	·	.70				



Common Name	Species Code	Scientific Name	Provincial Status (S Rank)	Global Status (G Rank)	SARO (MECP)	COSEWIC (Federal)	SWH Indicator Species	Highest Breeding Evidence
Galliformes								
Phasianinae	14/7-7-1		0.5	05			.,	50.11
Wild Turkey	WITU	Meleagris gallopavo	S5	G5			X	PO-H
Columbiformes								
Columbidae								
Mourning Dove	MODO	Zenaida macroura	S5	G5				PO-H
Trochilidae								
Ruby-throated Hummingbird	RTHU	Archilochus colubris	S5B	G5		1		PO-H
Suliformes								
Phalacrocoracidae								
Double-crested Cormorant	DCCO	Phalacrocorax auritus	S5B, S4N	G5	NAR	NAR		OB-X
Pelecaniformes								
Ardeidae								
Green Heron	GRHE	Butorides virescens	S4B	G5	·		Χ	OB-X
Great Blue Heron	GBHE	Ardea herodias	S5	G5	-			OB-X
Cathartiformes	<u> </u>							
Accipitriformes	<b>†</b>							
Accipitridae	<b>†</b>							
Red-tailed Hawk	RTHA	Buteo jamaicensis	S5	G5	NAR	NAR	X	PO-H
Piciformes								
Picidae								
Red-bellied Woodpecker	RBWO	Melanerpes carolinus	S5	G5				PO-H
Yellow-bellied Sapsucker	YBSA	Sphyrapicus varius	S5B, S3N	G5			X	PR-A
Downy Woodpecker	DOWO	Dryobates pubescens	S5	G5				PR-T
Hairy Woodpecker	HAWO	Dryobates villosus	S5	G5				PO-H
Northern Flicker	NOFL	Colaptes auratus	S5	G5				PO-H
Γ	1	T	1					
Passeriformes								
Tyrannidae	0.051		orn.	05				20.11
Great Crested Flycatcher	GCFL	Myiarchus crinitus	S5B	G5				PO-H
Eastern Kingbird Eastern Wood-Pewee	EAKI EAWP	Tyrannus tyrannus	S4B S4B	G5 G5	SC	SC	X	PO-H PR-T
Eastern Wood-Pewee	EAWP	Contopus virens	34D	G5	30	30	^	PR-1
Vireonidae								
Red-eyed Vireo	REVI	Vireo olivaceus	S5B	G5				PR-T
Corvidae								
Blue Jay	BLJA	Cyanocitta cristata	S5	G5				PR-T
American Crow	AMCR	Corvus brachyrhynchos	S5	G5				PO-H
Hirundinidae						+		
Tree Swallow	TRES	Tachycineta bicolor	S4S5B	G5				PO-H
Northern Rough-winged Swallow	NRWS	Stelgidopteryx serripennis	S4B	G5			X	OB-X
Paridae Black-capped Chickadee	BCCH	Poecile atricapillus	S5	G5				PR-T
	ВССП	Poeciie atricapilius	53	G5				PK-1
Sittidae								
Red-breasted Nuthatch	RBNU	Sitta canadensis	S5	G5			Х	РО-Н
White-breasted Nuthatch	WBNU	Sitta carolinensis	S5	G5		+		PR-A
Troglodytidae								
House Wren	HOWR	Troglodytes aedon	S5B	G5				PR-P
Turdidae								
Veery	VEER	Catharus fuscescens	S5B	G5			X	PR-A
Wood Thrush	WOTH	Hylocichla mustelina	S4B	G4	THR	THR	Х	PR-A
American Robin	AMRO	Turdus migratorius	S5	G5		1		PR-T
Fringillidae	-							
Purple Finch	PUFI	Haemorhous purpureus	S5	G5				PO-S
American Goldfinch	AMGO	Spinus tristis	S5	G5				PR-T

10249 Hunsden Sideroad Caledon, Ontario



Common Name	Species Code	Scientific Name	Provincial Status (S Rank)	Global Status (G Rank)	SARO (MECP)	COSEWIC (Federal)	SWH Indicator Species	Highest Breeding Evidence
Passerellidae								
Grasshopper Sparrow	GRSP	Ammodramus savannarum	S4B	G5	SC	SC	Х	PR-T
Chipping Sparrow	CHSP	Spizella passerina	S5B, S3N	G5				PR-A
Clay-colored Sparrow	CCSP	Spizella pallida	S4B	G5			X	PO-S
Field Sparrow	FISP	Spizella pusilla	S4B, S3N	G5			X	PR-T
Vesper Sparrow	VESP	Pooecetes gramineus	S4B	G5			Х	PR-T
Savannah Sparrow	SAVS	Passerculus sandwichensis	S5B, S3N	G5			Х	PO-S
Song Sparrow	SOSP	Melospiza melodia	S5	G5		•		PR-T
Eastern Towhee	EATO	Pipilo erythrophthalmus	S4B, S3N	G5			X	PO-S
Icteridae								
Eastern Meadowlark	EAME	Sturnella magna	S4B, S3N	G5	THR	THR		PR-T
Baltimore Oriole	BAOR	Icterus galbula	S4B	G5				CO-FY
Common Grackle	COGR	Quiscalus quiscula	S5	G5				PO-H
Parulidae								
Ovenbird	OVEN	Seiurus aurocapilla	S5B	G5			Х	PO-S
Mourning Warbler	MOWA	Geothlypis philadelphia	S5B	G5				PR-T
Common Yellowthroat	COYE	Geothlypis trichas	S5B	G5				PO-S
American Redstart	AMRE	Setophaga ruticilla	S5B	G5				PR-T
Chestnut-sided Warbler	CSWA	Setophaga pensylvanica	S5B	G5				PR-T
Black-throated Blue Warbler	BTBW	Setophaga caerulescens	S5B	G5			X	PR-A
Pine Warbler	PIWA	Setophaga pinus	S5B, S3N	G5				PR-T
Cardinalidae								
Scarlet Tanager	SCTA	Piranga olivacea	S5B	G5			X	PR-T
Northern Cardinal	NOCA	Cardinalis cardinalis	S5	G5				PO-S
Rose-breasted Grosbeak	RBGR	Pheucticus Iudovicianus	S5B	G5				CO-CF
Indigo Bunting	INBU	Passerina cyanea	S5B	G5				PR-A

51

Name:

Species Common Name and Scientific Chesser, R. T., K. J. Burns, C. Cicero, J. L. Dunn, A. W. Kratter, I. J. Lovette, P. C. Rasmussen, J. V. Remsen, Jr., D. F. Stotz, B. M. Winger, and K. Winker. 2018. Check-list of North American Birds (online). American Ornithological Society. Available online: http://checklist.aou.org/taxa

Species Code: Consistent with the American Ornithologists' Union. 2018. Species 4-Letter-Codes. Available online:

http://www.birdsontario.org/atlas/codes.jsp?lang=en&pg=species

**Highest Breeding Evidence:** Codes assigned for breeding evidence are consistent with the Ontario Breeding Bird Atlas (OBBA). 2018. Breeding Evidence

Codes. Available online: http://www.birdsontario.org/atlas/codes.jsp?lang=en&pg=breeding&sortorder=aou

Provincial ranks are from the Natural Heritage Information Centre; S1 (critically imperiled), S2 (imperlied), S3 (vulnerable), S ranks:

S4 (apparently secure), S5 (secure); ranks were updated using NHIC species list 2021. Available to download from:

https://www.ontario.ca/page/get-natural-heritage-information

Global ranks are from the Natural Heritage Information Centre; G1 (extremely rare), G2 (very rare), G3 (rare to G ranks:

uncommon), G4 (common), G5 (very common); ranks were updated using NHIC species list 2021. Available to download

from: https://www.ontario.ca/page/get-natural-heritage-information

SARO (MECP): Ontario Species at Risk as listed by the Committee on the Status of Species at Risk in Ontario (from Ontario Regulation

230/08 Species at Risk in Ontario website: https://www.ontario.ca/laws/regulation/080230/); END - Endangered; THR -

Threatened; SC - Special Concern; NAR - Not at Risk

COSEWIC: Assessed Species at Risk at the national level as listed by the Committee on the Status of Endangered Wildlife in Canada

(from COSEWIC: https://wildlife-species.canada.ca/species-risk-registry/sar/index/default\_e.cfm); END - Endangered, THR

- Threatened, SC - Special Concern, NAR - Not at Risk

SWH refers to Significant Wildlife Habitat as defined by the MNRF (2015) Significant Wildlife Habitat Criteria Schedules for **SWH Indicator Species:** 

Ecoregions 7E and 6E (as appropriate for the Subject Lands). SWH indicator species are identified in this table and any potential SWH is discussed in the text of this report. Available online: http://www.townofnemi.on.ca/wp-

content/uploads/2016/02/NEMI-OP-App-C-schedule-6e-jan-2015-access-ver-final-s.pdf



COMMON NAME	SCIENTIFIC NAME	Provincial Status (S RANK)	Global Status (G RANK)	SARO (MECP)	COSEWIC (Federal)	Local Status Halto	Local Status Hamilto n	Local Status TRCA	Regional Status Region of Waterloo	Local Status CVC	Niagara Region CA Status	SWH Indicato r Species 6E	SWH Indicator Species 7E
ODONATA	SCIENTIFIC NAME	RAINK)	KANK)	(MECP)	(reuerar)	- "		IKCA	waterioo	CVC	Status	OE.	/ [
Boreal Bluet	Enallagma boreale	S5	G5			HR	m						
Four-spotted Skimmer	Libellula quadrimaculata	S5	G5			HIK	- '''				С		
Common Whitetail	Plathemis lydia	S5	G5								C		
Common Winteran	riaciieiiiis iyala	33	0.5										
BUTTERFLIES													
Tawny-edged Skipper	Polites themistocles	S5	G5								С		
Hobomok Skipper	Poanes hobomok	S5	G5								_		
Black Swallowtail	Papilio polyxenes	S5	G5								Н		
Eastern Tiger Swallowtail	Papilio glaucus	S5	G5										
Cabbage White	Pieris rapae	SNA	G5										
Silvery Blue	Glaucopsyche lygdar	S5	G5								С		
Common Ringlet	Coenonympha tullia	S5	G5								R		
- Common rangisc	Goernerry ringing carrie	- 55	100										
BUMBLE BEES													
Two-spotted Bumble Bee	Bombus bimaculatus	S5	G5										
Common Eastern Bumble Bee	Bombus impatiens	S5	G5										
Northern Amber Bumble Bee	Bombus borealis	S5	G4G5										
AMBUTRIANG													
AMPHIBIANS	A	CF	C.F.								147		
American Toad	Anaxyrus americanus	S5	G5					L4	Х		W	X	Х
BIRDS											L		
Killdeer	Charadrius vociferus	S4B	G5					L4					
American Crow	Corvus brachyrhynchos	S5	G5					L5			С	-	
Bobolink	Dolichonyx oryzivorus	S4B	G5	THR	THR			L2			C		
BODOIIIK	Dollchonyx oryzivorus	340	GS	IIIK	IIIK			LZ					
MAMMALS													
Eastern Chipmunk	Tamias striatus	S5	G5					L4					
Eastern Gray Squirrel	Sciurus carolinensis	S5	G5					L5					
White-tailed Deer	Odocoileus virginianus	S5	G5					L4				Х	Х
SUMMARY													
JOHN AN I													
Total Odonata:		3											
Total Butterflies:		7											
Total Other Arthropods		3											
Total Amphibians:		1											
Total Reptiles:		0											
Total Birds:		0											
Total Breeding Birds:		3											
Total Mammals:		3											
SIGNIFICANT SPECIES													
Global:		0			<del> </del>								$\vdash$
National:		0											<del></del>
		0					-						<del></del>
Provincial:								1					<b></b>
Regional:		0											<b></b>
Local:							L	<u> </u>					



COMMON NAME	SCIENTIFIC NAME	Provincial Status (S RANK)	Global Status (G RANK)	SARO (MECP)	COSEWIC (Federal)	Local Status Halto n	Local Status Hamilto n	Local Status	Regional Status Region of Waterloo	Local Status CVC	Niagara Region CA Status	SWH Indicato r Species 6E	SWH Indicator Species 7E
COMMON NAME	SCIENTIFIC NAME	KAINK)	KANK)	(MECF)	(reuerar)			IRCA	Waterioo	CVC	Status	UE	/ =
Explanation of Status and Acronymns													
COSSARO: Committee on the Status of Species at Risk in Onta													
COSEWIC: Committee on the Status of Endangered Wildlife in	Canada												
S1: Critically Imperiled—Critically imperiled in the province (oft	,												
S2: Imperiled—Imperiled in the province, very few populations (													
S3: Vulnerable—Vulnerable in the province, relatively few popul	lations (often 80 or fewer)												
S4: Apparently Secure—Uncommon but not rare													
S5: Secure—Common, widespread, and abundant in the province	ce												
SX: Presumed extirpated													
SH: Possibly Extirpated (Historical)													
SNR: Unranked													
SU: Unrankable—Currently unrankable due to lack of information													
SNA: Not applicable—A conservation status rank is not applicable	ole because the species is not a suitable target	for conservation activities	es.										
S#S#: Range Rank—A numeric range rank (e.g., S2S3) is used	to indicate any range of uncertainty about the	status of the species											
S#B- Breeding status rank													
S#N- Non Breeding status rank													
?: Indicates uncertainty in the assigned rank													
G1: Extremely rare globally; usually fewer than 5 occurrences in	the overall range												
G1G2: Extremely rare to very rare globally													
G2: Very rare globally; usually between 5-10 occurrences in the	overall range												
G2G3: Very rare to uncommon globally													
G3: Rare to uncommon globally; usually between 20-100 occurr	ences												
G3G4: Rare to common globally													
G4: Common globally; usually more than 100 occurrences in the	e overall range												
G4G5: Common to very common globally													
G5: Very common globally; demonstrably secure													
GU: Status uncertain, often because of low search effort or cryp	tic nature of the species; more data needed.												
T: Denotes that the rank applies to a subspecies or variety													
Q: Denotes that the taxonomic status of the species, subspecies	s, or variety is questionable.												
END: Endangered													
THR: Threatened													
SC: Special Concern													
NAR: Not At Risk													
IND: Indeterminant, insufficient information to assign status													
DD: Data Deficient													
6: Rare in Site Region 6													
7: Rare in Site Region 7													
Area: Minimum patch size for area-sensitive species (ha)				-									1
H- highly significant in Hamilton Region (i.e. rare)													
m- moderately significant in Hamilton Region (i.e. uncommon)													
L1- extremely rare locally (Toronto Region)													
L2- very rare locally (Toronto Region)													
L3- rare to uncommon locally (Toronto Region)													
HR- rare in Halton Region, highly significant													
HU- uncommon in Halton Region, moderately significant													
REFERENCES													
COSSARO Status													
Endangered Species Act, 2007 (Bill 184). Species at Risk in On	ntario List (O. Reg. 230/08). Accessed October	7, 2016.											
	- ·												



COMMON NAME	SCIENTIFIC NAME	Provincial Status (S RANK)	Global Status (G RANK)	SARO (MECP)	COSEWIC (Federal)	Local Status Halto n	Local Status Hamilto n	•	Local Status	Niagara Region CA Status	SWH Indicato r Species 6E	SWH Indicator Species 7E
COSEWIC Status												
COSEWIC. 2016. Canadian Species at Risk. Committee on the	he Status of Endangered Wildlife in Canada.											
Local Status												
Dwyer, Jill K. 2003. Nature Counts Project Hamilton Natural A	reas Inventory 2003. Species Checklists. Ham	ilton Naturalists Club.										
Halton Natural Areas Inventory. 2006. Volume 2 Species Ched	klists (ISBN 0-9732488-7-4).											
Region of Waterloo. 1996. Regionally Significant Breeding Bi	rds.											
Toronto and Region Conservation Authority (TRCA). 2016. Re	vised Fauna Scores and Ranks, February 2016											
Hamilton Conservation Authority (HCA). 2014. Hamilton Natur	al Areas Inventory Project (3rd Edition).											
Significant Wildlife Habitat (SWH) Indicator Species												
Ministry of Natural Resources and Forestry (MNRF). 2015. Sig Available at: https://dr6j45jk9xcmk.cloudfront.net/documents/4												
Ministry of Natural Resources and Forestry (MNRF). 2015. Sig Available at: https://dr6j45jk9xcmk.cloudfront.net/documents/4												
Natural Heritage Information Center (NHIC). 2016. Onatrio Sp	ecies List: All Species.											



Table 5: Significant Wildlife Habitat Assessment (6E)

SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?
1. SEASONAL CONCENTRATI	ON AREAS			
Waterfowl Stopover and Staging Areas (terrestrial)	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Waterfowl Stopover and Staging Areas (aquatic)	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Shorebird Migratory Stopover Areas	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Raptor Wintering Areas	No – suitable upland ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Bat Hibernacula	No – Cave and Crevice communities are absent from the Subject Lands.	N/A	No	No – SWH type is not present
Bat Maternity Colonies	Yes – Forested (FOD) vegetation communities are present within the Subject Lands.	Additional studies would be required to confirm if habitat conditions are met.	No – Avoidance of woodland communities is recommended.	Yes – SWH type may be present; treated as Candidate SWH
Turtle Wintering Areas	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Reptile Hibernacula	Yes – ecosites are present on the Subject Lands.	A rock pile feature was identified in the FOD6-5.	No – Avoidance of woodland communities is recommended.	Yes – SWH type may be present; treated as Candidate SWH



#### Table 5: Significant Wildlife Habitat Assessment (6E)

SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?
Colonial Bird Nesting Sites (bank/cliff)	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Colonial Bird Nesting Sites (tree/shrubs)	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Colonial Bird Nesting Sites (ground)	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
Migratory Butterfly Stopover Areas	No – suitable ecosites are not present within the Subject Lands.	No – The Subject Lands are located greater than 5 km away from Lake Ontario.	No	No – SWH type is not present
Migratory Landbird Stopover Areas	Yes – FO vegetation communities are identified within the Subject Lands.	No – The Subject Lands are located greater than 5 km away from Lake Ontario.	No	No – SWH type is not present
Deer Yarding Areas	No – Mapping from the MNRF LIO database did not depict any deer yarding areas on or adjacent to the Subject Lands.	N/A	No	No – SWH type is not present
Deer Winter Congregation Areas	No – Mapping from the MNRF LIO database did not depict any deer wintering areas on or adjacent to the Subject Lands.	N/A	No	No – SWH type is not present



#### Table 5: Significant Wildlife Habitat Assessment (6E)

SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?	
2. RARE VEGETATION COMMUNITIES OR SPECIALIZED HABITAT FOR WILDLIFE					
2a. Rare Vegetation Communitie	es				
Rare Vegetation Types (cliffs, talus slopes, sand barrens, alvars, old-growth forests, savannahs, and tallgrass prairies)	No – None identified through the background information review or site reconnaissance.	N/A	No	No – SWH type is not present	
Other Rare Vegetation Types (S1 to S3 communities)	No – None identified thought the background information review or site reconnaissance.	N/A	No	No – SWH type is not present	
2b. Specialized Wildlife Habitat					
Waterfowl Nesting Area	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present	
Bald Eagle and Osprey Habitats	Yes – FO ecosites are present within the Subject Lands.	No permanent watercourses are present within the forested communities and no large stick nests were observed within the Subject Lands	Yes – breeding bird surveys were completed.	No – SWH type is not present.	
Woodland Raptor Nesting Habitat	Yes – FO ecosites are present within the Subject Lands.	Yes – The south-most FO feature is a part of a large contiguous forest system and would meet the required woodland size and interior habitat size is not achieved.	Yes – breeding bird surveys were completed; however, none of the target raptors species were recorded.	No – SWH type is not present	



Table 5: Significant Wildlife Habitat Assessment (6E)

SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?	
Turtle Nesting Areas	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present	
Seeps and Springs	Yes – Forested ecosites are present within the Subject Lands.	No evidence of seeps or springs was identified within the Subject Lands.	No	No – SWH type is not present	
Woodland Amphibian Breeding Habitats (within or < 120m from woodland)	Yes – Forested ecosites with are present within the Subject Lands.	No vernal pools were identified within the Subject Lands.	No	No – SWH type is not present	
Wetland Amphibian Breeding Habitats (wetland >120m from woodland)	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present	
Woodland Area-Sensitive Bird Breeding Habitat	Yes – FO ecosites are present within the Subject Lands.	Yes – The south-most FO feature is a part of a large contiguous forest system and would meet the required woodland size and interior habitat size is not achieved.	Yes – breeding bird surveys were completed and several of the target bird species were recorded as probable and possible breeders. This includes: Veery, Black-throated Blue Warbler and Scarlet Tanager	Yes – SWH type is present in the southern most Forest Community since three of the listed wildlife species are probable or possible breeders within the Subject Lands.	



Table 5: Significant Wildlife Habitat Assessment (6E)

SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?		
3. SPECIES OF CONSERVATION	3. SPECIES OF CONSERVATION CONCERN					
Marsh Bird Breeding Habitat	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present		
Open Country Bird Breeding Habitat	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present		
Shrub/Early Successional Bird Breeding Habitat	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present		
Terrestrial Crayfish	No – suitable ecosites are not present within the Subject Lands.	N/A	No, however no crayfish chimneys were identified during the field investigations.	No – SWH type is not present		
Special Concern and Rare Wildlife Species (based on the Secondary Source Review – Section 3.0)						
(i) Common Nighthawk - SC	N/A	No – the preferred habitat types of the species (i.e., logged or burned-over areas, forest clearings, rock barrens, peat bogs, lakeshores, and mine tailing) are not present within the Subject Lands	No	No – SWH type is not present		



#### Table 5: Significant Wildlife Habitat Assessment (6E)

SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?
(ii) Eastern Wood-Pewee - SC	N/A	Possibly – Forested ecosites are present within the Subject Lands.	Yes – Breeding bird surveys were completed, and the species was recorded in both Forested communities within the Subject Lands.	Yes – SWH type is present in both forests within the Subject Lands.
(iii) Grasshopper Sparrow - SC	N/A	No – Suitable grasslands habitats are not present within the Subject Lands	Yes – Breeding bird surveys were completed. This species was recorded outside of the Subject Lands but not within the Subject Lands.	No – SWH type is not present
(iv) Golden-winged Warbler - SC	N/A	No – While field edges, a preferred habitat type of the species, is present within the Subject Lands; the Subject Lands are not located within the known occurrence range of the species (MECP 2021).	Yes – Breeding bird surveys were completed, and the species was not present.	No – SWH type is not present
(v) Wood Thrush -SC	N/A	Possibly – Forested ecosites are present within the Subject Lands.	Yes – Breeding bird surveys were completed, and the species was recorded in both Forested communities within the Subject Lands.	Yes – SWH type is present in both forests within the Subject Lands.



Table 5: Significant Wildlife Habitat Assessment (6E)

SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?
(vi) Common Gallinule – S3B	N/A	No – Suitable wetlands are not present within the Subject Lands.	Yes – Breeding bird surveys were completed, and the species was not present.	No – SWH type is not present
(vii) Purple Martin – S3B	N/A	No – This species almost exclusively nests in artificial roosting boxes. No roosting boxes were present within the Subject Lands; therefore, the species is not expected to be present.	Yes – Breeding bird surveys were completed, and the species was not present.	No – SWH type is not present
(viii) Blue-winged Teal -S3B	N/A	No – Suitable wetlands are not present within the Subject Lands.	Yes – Breeding bird surveys were completed, and the species was not present.	No – SWH type is not present
(ix)Monarch - SC	No – suitable ecosites are not present within the Subject Lands.	N/A	No	No – SWH type is not present
(x) Lilypad Clubtail S3	N/A	No – suitable aquatic habitats are not present within the Subject Lands.	No	No – SWH type is not present
(xi)Snapping Turtle - SC	N/A	No – Suitable wetlands are not present within the Subject Lands.	No	No – SWH type is not present



Table 5: Significant Wildlife Habitat Assessment (6E)


SIGNIFICANT WILDLIFE HABITAT (SWH) TYPE	ELC ECOSITE(S) PRESENT	HABITAT CRITERIA MET	TARGETED FIELD STUDIES REQUIRED	POTENTIAL FOR SWH TYPE PRESENCE?
(xii) Nerveless Kuhlenberg's Sedge S3	NA	Possibly, this species prefers dry grasslands, open forests and commonly occurs on sand and acidic soil.	Yes – a botanical inventory was completed, and the species was not present.	No – SWH type is not present
4. ANIMAL MOVEMENT CORRIDORS				
Amphibian Movement Corridors	N/A	No	No	No – SWH type is not present

Ministry of Environment Conservation and Parks (MECP). 2021. Golden-winged Warbler Website. Available Online at <a href="https://www.ontario.ca/page/golden-winged-warbler">https://www.ontario.ca/page/golden-winged-warbler</a>

# Appendix C

**Draft Plan** 





# DRAFT PLAN OF SUBDIVISION SUZANNE WILSON FILE # 21T-22004C

PART OF LOTS 25 AND 26, CONCESSION 9 (ALBION)
PART OF ROAD ALLOWANCE BETWEEN PARTS 25 AND
26, CONCESSION 9 (ALBION)
TOWN OF CALEDON
REGIONAL MUNICIPALITY OF PEEL

## **OWNERS CERTIFICATE**

I HEREBY AUTHORIZE GLEN SCHNARR & ASSOCIATES INC. TO PREPARE AND SUBMIT THIS DRAFT PLAN OF SUBDIVISION TO THE TOWN OF CALEDON FOR APPROVAL.

SIGNED: Suzanne Wilson

DATE: June 29, 2022

# **SURVEYORS CERTIFICATE**

I HEREBY CERTIFY THAT THE BOUNDARIES OF THE LANDS TO BE SUBDIVIDED AS SHOWN ON THIS PLAN AND THEIR RELATIONSHIP TO ADJACENT LANDS ARE CORRECTLY AND ACCURATELY SHOWN.

SIGNED: 4. Stedmill OLS

DATE: <u>July 6, 2022</u>

GRANT T. STIDWILL, O.L.S.

J.D. BARNES LIMITED

401 WHEELABRATOR WAY, SUITE A
MILTON, ON

MILTON, ON TEL.: (905) 875-9955 WEB: www.jdbarnes.com

## ADDITIONAL INFORMATION

(UNDER SECTION 51(17) OF THE PLANNING ACT) INFORMATION REQUIRED BY CLAUSES A,B,C,D,E,F,G,J & L ARE SHOWN ON THE DRAFT AND KEY PLANS.

H) MUNICIPAL AND PIPED WATER TO BE PROVIDED

I) SANDY LOAM AND CLAY LOAM

K) SERVICED BY SEPTIC SYSTEMS.

# LAND USE SCHEDULE

LAND USE	LOTS / BLOCKS	AREA (ha)	AREA (ac)	UNITS
ESTATE LOTS	1-11	6.303	15.575	11
BONUS ESTATE LOTS	12 & 13	1.362	3.365	2
EX. WILSON DWELLING PARCEL, EPA2-ORM	14	1.214	2.999	
NHS	15	7.974	19.703	
NHS BUFFER	16	2.307	5.701	
OPEN SPACE	17 & 18	0.151	0.373	
0.3m RESERVE	19 & 20	0.00	0.000	
18.0m LOCAL R.O.W. (LENGTH: 566m)		1.064	2.630	
TOTAL	20	20.374	50.346	13

# NOTES

0.151093492328

-PAVEMENT ILLUSTRATION IS DIAGRAMMATIC
-EXISTING RESIDENCE TO REMAIN
-DAYLIGHT ROUNDINGS 5m UNLESS OTHERWISE NOTED

-DRIPLINE AS STAKED BY THE CONSERVATION AUTHORITY DATED AUGUST 09 2023 -STRUCTURAL ENVELOPE MINIMUM 30M FROM NHS BLOCK 15



SCALE 1:1500 (24 x 36) MARCH 11, 2024

